

UTILITY  
PATENT APPLICATION  
TRANSMITTAL

Only for new nonprovisional applications under 37 CFR 1.53(b)

Attorney Docket No.	MIT-087
First Named Inventor	Rosenberg et al.
Title	HEPARAN SULFATE D-GLUCOSAMINYL 3-O-SULFOTRANSFERASES, AND USES THEREFOR

## APPLICATION ELEMENTS

ADDRESS TO: **Box Patent Application**  
Assistant Commissioner for Patents  
Washington, D.C. 20231

1. ☒ Fee Transmittal Form
2. ☒ Specification and Drawings [77 Total Pages]  
- Specification & listed references - (64 pages)  
- Claims - (9 pages)  
- Abstract - (1 page)  
- Sheets of Drawings - (3 sheets)  
☐ Formal  
☒ Informal

## ACCOMPANYING APPLICATION PARTS

7. ☐ Associate Power of Attorney
8. ☐ English Translation Document (if applicable)
9. ☐ Information Disclosure Statement (IDS)/PTO-1449  
☐ Copies of IDS Citations
10. ☐ Preliminary Amendment (\_\_\_ pages)  
☐ Drawings [Total Sheets \_\_\_]  
☐ Letter to Official Draftsperson Including Drawings [Total Pages \_\_\_]

3. ☐ Oath or Declaration [Total Pages \_\_\_]  
a. ☐ Newly executed (original)  
b. ☐ Copy from a prior application (37 CFR 1.63(d))  
(for continuation/divisional with Box 17 completed)  
[Note Box 4 below]

11. ☒ Return Receipt Postcard

4. ☐ Incorporation by Reference (usable if Box 3b is checked)  
The entire Disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 3b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

12. ☒ Small Entity Statement (1 page)  
☐ Statements filed in prior application, (Status still proper and desired)
13. ☐ Certified Copy of Priority Document(s)

5. ☐ Microfiche Computer Program (Appendix)

14. ☐ Deletion of Inventor(s)  
Signed statement attached deleting inventor(s) named in the prior application.

6. ☒ Nucleotide and/or Amino Acid Sequence Submission  
☒ Computer Readable Copy  
☒ Paper Copy (30 pgs.) (identical to computer copy)  
☒ Statement verifying identity of above copies

15. ☐ Patent Application Data Entry Form
16. ☐ Other:

17. ☒ Priority - 35 U.S.C. 120

This application claims benefit of priority of International Patent Application Serial No. PCT/US98/22597, with an international filing date of October 23, 1998, which claims priority to U.S. Provisional Patent Application Serial No. 60/062,762 filed on October 24, 1997, and U.S. Provisional Patent Application Serial No. 60/065,437, filed on October 31, 1997.

Priority to International Patent Application Serial No. PCT/US98/22597 is claimed under 35 U.S.C. 120.

18. ☒ Priority - 35 U.S.C. 119

This application claims benefit of priority of International Patent Application Serial No. PCT/US98/22597, with an international filing date of October 23, 1998, which claims priority to U.S. Provisional Patent Application Serial No. 60/062,762 filed on October 24, 1997, and U.S. Provisional Patent Application Serial No. 60/065,437, filed on October 31, 1997.


Priority to U.S. Provisional Patent Application Serial No. 60/062,762 and U.S. Provisional Patent Application Serial No. 60/065,437 is claimed under 35 U.S.C. 119(e).

- ☐ The certified copy has been filed in prior U.S. application Serial No. \_\_\_\_/\_\_\_\_ on \_\_\_\_.
- ☐ The certified copy will follow.

## CORRESPONDENCE ADDRESS

Direct all correspondence to: Patent Administrator  
Testa, Hurwitz & Thibault, LLP  
High Street Tower  
125 High Street  
Boston, MA 02110  
Tel. No.: (617) 248-7000  
Fax No.: (617) 248-7100

## SIGNATURE BLOCK

Respectfully submitted,  
  
Maureen A. Bresnahan  
Attorney for Applicants  
Testa, Hurwitz & Thibault, LLP  
High Street Tower  
125 High Street  
Boston, MA 02110

Date: April 24, 2000  
Reg. No. 44,559  
Tel. No.: (617) 248-7103  
Fax No.: (617) 248-7100

EXPRESS MAIL NO.: EL302647095US

**STATEMENT CLAIMING SMALL ENTITY STATUS  
(37 CFR 1.9(f) & 1.27(d)) - NONPROFIT ORGANIZATION**

Attorney Docket No. MIT-087

Applicants: Robert D. Rosenberg, Nicholas W. Shwornak, Jian Liu, Linda M.S. Fritze, John J. Schwartz and Lijuan ZhangApplication or Patent No.: Not yet assignedFiled: Filed herewithTitle: HEPARAN SULFATE D-GLUCOSAMINYL 3-O-SULFOTRANSFERASES AND USES THEREFOR

I hereby state that I am an official empowered to act on behalf of the nonprofit organization identified below:  
NAME OF NONPROFIT ORGANIZATION Massachusetts Institute of Technology  
ADDRESS OF NONPROFIT ORGANIZATION 77 Massachusetts Avenue, Cambridge, MA 02139

**TYPE OF NONPROFIT ORGANIZATION:**

- ☒ UNIVERSITY OR OTHER INSTITUTION OF HIGHER EDUCATION
- ☐ TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 U.S.C. 501(a) and 501(c)(3))
- ☐ NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA  
(NAME OF STATE \_\_\_\_\_)  
(CITATION OF STATUTE \_\_\_\_\_)
- ☐ WOULD QUALIFY AS TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 U.S.C. 501(a) and 501(c)(3)) IF LOCATED IN THE UNITED STATES OF AMERICA
- ☐ WOULD QUALIFY AS NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA IF LOCATED IN THE UNITED STATES OF AMERICA  
(NAME OF STATE \_\_\_\_\_)  
(CITATION OF STATUTE \_\_\_\_\_)

I hereby state that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9(e) for purposes of paying reduced fees to the United States Patent and Trademark Office regarding the invention described in:

- ☒ the specification filed herewith with title as listed above.
- ☐ the application Serial No. \_\_\_\_\_, filed \_\_\_\_\_
- ☐ the patent identified above.

I hereby state that the rights under contract or law have been conveyed to and remain with the nonprofit organization regarding the above identified invention. If the rights held by the nonprofit organization are not exclusive, each individual, concern, or organization having rights in the invention must file separate statements as to their status as small entities and that no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(e) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

Each person, concern, or organization having any rights in the invention is listed below:

- ☒ no such person, concern, or organization exists.
- ☐ each such person, concern, or organization is listed below.

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

NAME OF PERSON SIGNING Jarmila Z. HrbekTITLE IN ORGANIZATION OF PERSON SIGNING Patent AdministratorADDRESS OF PERSON SIGNING Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139SIGNATURE Jarmila Z. Hrbek DATE April 21, 2000

**HEPARAN SULFATE D-GLUCOSAMINYL 3-O-SULFOTRANSFERASES,  
AND USES THEREFOR**

Cross-Reference to Related Application

This application claims benefit of priority of International Patent Application  
Serial No. PCT/US98/22597, with an international filing date of October 23, 1998,  
which claims priority to U.S. Provisional Patent Application Serial No. 60/062,762,  
5 filed on October 24, 1997, and U.S. Provisional Patent Application Serial No.  
60/065,437, filed on October 31, 1997.

Field of the Invention

The present invention is related to the field of biochemistry and molecular  
10 biology, and in particular to the field of enzymology and heparan sulfate biosynthesis.

Background of the Invention

The serine proteases of the intrinsic blood coagulation cascade are slowly  
neutralized by antithrombin (AT) (reviewed in (1)). This inhibition is secondary to  
the generation of 1:1 enzyme-AT complexes whose formation is dramatically  
15 enhanced by the mast cell product, heparin (2). Damus *et al.* (3) hypothesized that  
endothelial cell surface heparan sulfate proteoglycans (HSPGs) function in a similar  
fashion to accelerate coagulation enzyme inactivation by AT, and therefore are  
responsible for the non-thrombogenic properties of blood vessels. It was initially  
demonstrated that perfusion of the hindlimbs of normal rodents and rodents deficient  
20 in mast cells with purified thrombin (T) and AT leads to a greatly elevated rate of T-  
AT complex formation and that the enzyme heparitinase as well as the natural heparin  
antagonist platelet factor 4 suppress the above acceleration (4, 5). It was subsequently  
showed that cultured cloned bovine macrovascular and rodent microvascular  
endothelial cells synthesize both anticoagulant HSPG (HSPG<sup>act</sup>) as well as  
25 nonanticoagulant HSPG (HSPG<sup>inact</sup>) (6-8). HSPG<sup>act</sup> bear glycosaminoglycan (GAG)  
chains that bind tightly to AT and accelerate T-AT complex generation (6-8).

The biosynthesis of HSPG<sup>act</sup> requires generation of a core protein, assembly of a linkage region of four neutral sugars on specific serine attachment sites of the core protein, elongation of a GAG backbone composed of alternating *N*-acetylglucosamine and glucuronic acid residues, and modification of this homogenous copolymer by partial *N*-deacetylation with coupled *N*-sulfation of glucosamine residues, partial epimerization of glucuronic acid to iduronic acid residues, partial 2-*O*-sulfation of uronic acid residues, and partial 6-*O*-sulfation and partial 3-*O*-sulfation of glucosamine residues (reviewed in 9)). This multienzyme pathway generates HSPG<sup>act</sup> with regions of defined structure that contain the primary AT binding domain sequence found in anticoagulant heparin: uronic acid→glucosamine (*N*-acetyl/*N*-sulfate) 6-*O*-sulfate→glucuronic acid→glucosamine *N*-sulfate 3-*O*-sulfate (6-*O*-sulfate)→iduronic acid 2-*O*-sulfate→glucosamine *N*-sulfate 6-*O*-sulfate (10-17). These reactions also produce HSPG<sup>inact</sup> with regions of varying monosaccharide sequence that lack the primary AT -binding domain. The structure-function relationships of the AT binding domain have been elucidated with heparin/heparan sulfate oligosaccharides in association with fast reaction kinetics and equilibrium binding assays. The 6-*O*-sulfate group on residue 2 and the 3-*O*-sulfate group on residue 4 function in a thermodynamically linked fashion to supply half of the binding energy for interaction with AT, and trigger a conformational event that accelerates neutralization of specific coagulation proteases (11, 12). The amino and ester sulfate groups at residues 5 and 6, as well as carboxyl groups at other sites, provide the other half of the binding energy for interaction with protease inhibitor (10, 11). Furthermore, monosaccharide sequences outside the primary AT binding domain are essential in facilitating inhibition of coagulation proteases other than factor Xa (18, 19).

During the past eight years, several biosynthetic enzymes that generate HSPG<sup>act</sup> and HSPG<sup>inact</sup> have been purified. These proteins include an *N*-acetylglucosamine/glucuronic acid copolymerase (20), *N*-deacetylase/*N*-sulfotransferases (NST-1 and NST-2) (21, 22), a glucuronic acid/iduronic acid epimerase (23), an iduronic acid/glucuronic acid 2-*O*-sulfotransferase (2-OST) (24), a glucosamine 6-*O*-sulfotransferase (6-OST) (25) and a glucosamine 3-*O*-sulfotransferase (3-OST) (26, 35). However, the only enzymes that have also been



00440-232560

molecularly cloned are two structurally and functionally distinct isoforms of *N*-deacetylase/*N*-sulfotransferase (NST-1 from liver and NST-2 from mastocytoma) (27-31), and the 2-OST and epimerase. The above enzymes must function in a coordinated manner to produce the AT binding domain because the abundance of this sequence is much greater than predicted from a random assembly of constituents (32). The postulated regulatory mechanism must direct the biosynthetic enzymes to carry out the appropriate sequence of epimerization/sulfation reactions to generate the AT binding domain (33, 34).

#### Summary of the Invention

10           The present invention depends, in part, upon the identification and molecular cloning of novel genes encoding mammalian heparan sulfate D-glucosaminyl 3-O-sulfotransferases (3-OSTs). In particular, as disclosed herein, the present invention provides nucleic acid (SEQ ID NO: 1) and amino acid (SEQ ID NO: 2) sequences for murine 3-OST-1; nucleic acid (SEQ ID NO: 3) and amino acid (SEQ ID NO: 4) sequences for human 3-OST-1; nucleic acid (SEQ ID NO: 5) and amino acid (SEQ ID NO: 6) sequences for human 3-OST-2; nucleic acid (SEQ ID NO: 7) and amino acid (SEQ ID NO: 8) sequences for human 3-OST-3A; nucleic acid (SEQ ID NO: 9) and amino acid (SEQ ID NO: 10) sequences for human 3-OST-3B; and nucleic acid (SEQ ID NO: 11) and amino acid (SEQ ID NO: 12) sequences for human 3-OST-4. In addition, the invention provides amino acid (SEQ ID NO: 15) sequences for a *C. elegans* homologue, ce3-OST.

25           Thus, in one aspect, the present invention provides isolated nucleic acids encoding at least a functional fragment of a 3-OST protein. In preferred embodiments, the nucleic acid encodes a 3-OST protein comprising a mature murine or human 3-OST-1. In other embodiments, the nucleic acid encodes a 3-OST protein selected from 3-OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4, and ce3-OST. In other preferred embodiments, the nucleic acid encodes a 3-O-sulfotransferase domain of a 3-OST protein selected from 3-OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4, and ce3-OST. In particular embodiments, the nucleic acid comprises a nucleotide sequence selected from nucleotide sequences within: (a) SEQ ID NO: 1; (b) SEQ ID NO: 3; (c) SEQ ID NO: 5; (d) SEQ ID NO: 7; (e) SEQ ID NO: 9; (f) SEQ ID

30

NO: 11; (g) a sequence having at least 60% nucleotide sequence identity with at least one of (a)-(f) and encoding a functional fragment having sequence-specific HS binding affinity or 3-O-sulfotransferase activity; and (h) a sequence differing from a sequence of (a)-(g) only by the substitution of synonymous codons. In other particular  
5 embodiments, the present invention provides an isolated nucleic acid encoding a polypeptide selected from: (a) residues 21-52, 260-269, 250-276, 53-311, or 21-307 of SEQ ID NO: 2; (b) residues 21-48, 256-265, 246-272, 49-307, or 21-303 of SEQ ID NO: 4; (c) residues 42-109, 313-325, 303-332, or 110-367 of SEQ ID NO: 6; (d) residues 44-147, 351-363, 341-370, or 148-406 of SEQ ID NO: 8; (e) residues 66-  
10 132, 336-348, 326-355, or 133-390 of SEQ ID NO: 10; (f) residues 396-408, 386-415, or 207-456 of SEQ ID NO: 12; (g) residues 240-250, 230-257, 23-291 of SEQ ID NO: 15, (h) a sequence having at least 60% amino acid sequence similarity with at least one of (a)-(g) and encoding a functional fragment having sequence-specific HS binding affinity or 3-O-sulfotransferase activity; and (i) a sequence comprising a  
15 chimera of at least two of sequences (a)-(h).

In another aspect, the present invention provides isolated nucleic acids comprising at least 16 consecutive nucleotides of a nucleotide sequence selected from SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, and SEQ ID NO: 11.

20 In another aspect, the present invention provides for cells and cell lines transformed with the nucleic acids of the present invention. Thus, the invention provides host cells transformed with any of the above-described nucleic acids. The transformed host cells may be bacterial, yeast, or insect cells. Preferably, however, the host cells are mammalian cells, including endothelial cells, mast cells, fibroblasts,  
25 hybridomas, oocytes, and embryonic stem cells. Examples of preferred mammalian cells include COS-7 cells, murine primary cardiac microvascular endothelial cells (CME), murine mast cell line C57.1, primary human endothelial cells of umbilical vein (HUVEC), F9 embryonal carcinoma cells, rat fat pad endothelial cells (RFPEC), L cells (e.g., murine LTA *tk* cells), and cells derived from the transgenic animals of  
30 the invention. The transformed host cells may also be fetal cells, embryonic stem cells, zygotes, gametes, or germ line cells. Transformed embryonic stem cells, zygotes, gametes, and germ line cells, as well as other mammalian cells, may be used

to produce transgenic animals in which the expression of 3-OST genes have been altered (e.g., knock-outs, enhanced expression, ectopic expression).

In another aspect, the present invention provides substantially pure protein preparations comprising at least a functional fragment of a 3-OST protein. Thus, in

one embodiment, the present invention provides a substantially pure protein

preparation comprising mature murine 3-OST-1 or mature human 3-OST-1. In

another embodiment, the 3-OST protein is selected from the group consisting of 3-

OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4, and ce3-OST. In another

embodiment, the fragment comprises a 3-O-sulfotransferase domain of a 3-OST

protein selected from the group consisting of 3-OST-1, 3-OST-2, 3-OST-3A, 3-OST-

3B, 3-OST-4, and ce3-OST. In particular embodiments, the present invention

provides a substantially pure protein preparation in which the 3-OST protein

comprises an amino acid sequence selected from: (a) SEQ ID NO: 2; (b) SEQ ID

NO: 4; (c) SEQ ID NO: 6; (d) SEQ ID NO: 8; (e) SEQ ID NO: 10; (f) SEQ ID

NO: 12; (g) SEQ ID NO 15; and (h) a sequence having at least 60% amino acid

similarity with at least one of (a)-(g) and having sequence-specific HS binding affinity

or 3-O-sulfotransferase activity. In other particular embodiments, the present

invention provides a substantially pure protein preparation in which the 3-OST protein

comprises an amino acid sequence selected from: (a) residues 21-52, 260-269, 250-

276, 53-311, or 21-307 of SEQ ID NO: 2; (b) residues 21-48, 256-265, 246-272, 49-

307, or 21-303 of SEQ ID NO: 4; (c) residues 42-109, 313-325, 303-332, or 110-367

of SEQ ID NO: 6; (d) residues 44-147, 351-363, 341-370, or 148-406 of SEQ ID

NO: 8; (e) residues 66-132, 336-348, 326-355, or 133-390 of SEQ ID NO: 10; (f)

residues 396-408, 386-415, or 207-456 of SEQ ID NO: 12; (g) residues 240-250,

230-257, 23-291 of SEQ ID NO: 15; (h) a sequence having at least 60% amino acid

sequence similarity with at least one of (a)-(g) and encoding a functional fragment

having sequence-specific HS binding affinity or 3-O-sulfotransferase activity; and (i)

a sequence comprising a chimera of at least two of sequences (a)-(h).

In another aspect, the present invention provides for antibodies and methods

for making antibodies which selectively bind with the 3-OST proteins. These

antibodies include monoclonal and polyclonal antibodies, as well as functional

antibody fragments such as F(ab) and Fc.

In another aspect, the present invention provides for methods for producing the above-described proteins. Thus, in one set of embodiments, the isolated nucleic acids of the invention may be used to transform host cells or create transgenic animals which express the proteins of the invention. The proteins may then be substantially purified from the cells or animals by standard methods. Alternatively, the isolated nucleic acids of the invention may be used in cell-free *in vitro* translation systems to produce the proteins of the invention.

In another aspect, the present invention provides methods for 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides by contacting the preparation with at least a 3-O-sulfotransferase domain of a 3-OST protein in the presence of a sulfate donor under conditions which permit sulfation of the residues, and wherein the 3-OST protein is selected from 3-OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4, and ce3-OST proteins, as well as conservative substitution variants and/or chimeras thereof. In particular embodiments, the present invention provides methods for 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides in which the polysaccharides include a polysaccharide sequence of GlcA→GlcNS ±6S. These methods comprise contacting the GlcA→GlcNS ±6S-containing polysaccharide preparation with a 3-OST-1 protein in the presence of a sulfate donor under conditions which permit the 3-OST-1 to convert the GlcA→GlcNS ±6S sequence to GlcA→GlcNS 3S ±6S. In particular embodiments, the GlcA→GlcNS ±6S sequence comprises a part of an HS<sup>act</sup> precursor sequence (i.e., IdoA→GlcNAc 6S→GlcA→GlcNS ±6S→IdoA 2S→GlcNS 6S or IdoA→GlcNS 6S→GlcA→GlcNS ±6S→IdoA 2S→GlcNS 6S) or a part of an HS<sup>inact</sup> precursor sequence (i.e., IdoA→GlcNAc→GlcA→GlcNS ±6S→IdoA 2S→GlcNS 6S; IdoA→GlcNS→GlcA→GlcNS ±6S→IdoA 2S→GlcNS 6S; IdoA→GlcNAc 6S→GlcA→GlcNS ±6S→IdoA 2S→GlcNS; or IdoA→GlcNS 6S→GlcA→GlcNS ±6S→IdoA 2S→GlcNS). Conversion of the HS<sup>act</sup> precursor pool to HS<sup>act</sup> increases the fraction with AT-binding activity and is particularly useful in the production of anticoagulant heparan sulfate products. Thus, in another embodiment, the present invention provides for means of enriching the AT-binding fraction of a heparan

sulfate pool by contacting the polysaccharide preparation with 3-OST-1 protein in the presence of a sulfate donor under conditions which permit the 3-OST HS<sup>act</sup> conversion activity. The 3-OST-1 protein for use in these methods is selected from murine 3-OST-1, human 3-OST-1, mature murine 3-OST-1, mature human 3-OST-1, a functional fragment of a 3-OST-1 having 3-O-sulfotransferase activity, a conservative substitution variant of 3-OST-1 having 3-O-sulfotransferase activity, and a chimeric 3-OST-1 having 3-O-sulfotransferase activity. In preferred embodiments, the sulfate donor is 3'-phospho-adenosine 5'-phosphosulfate (PAPS).

Similarly, the present invention provides methods for 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides by contacting the preparation with at least a 3-O-sulfotransferase domain of a 3-OST protein in the presence of a sulfate donor under conditions which permit sulfation of the residues, and wherein the 3-OST protein is selected from 3-OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4, ce3-OST and conservative substitution variants or chimeras thereof. In particular embodiments, the present invention provides methods for 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides in which the polysaccharides include a polysaccharide sequence of GlcA 2S→GlcNS. These methods comprise contacting the GlcA 2S→GlcNS-containing polysaccharide preparation with a 3-OST-2 protein in the presence of a sulfate donor under conditions which permit the 3-OST-2 protein to convert the GlcA 2S→GlcNS sequence to GlcA 2S→GlcNS 3S. In particular embodiments, the GlcA 2S→GlcNS sequence comprises a part of a GlcNS→GlcA 2S→GlcNS sequence. In other particular embodiments, the present invention provides methods for 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides in which the polysaccharides include a polysaccharide sequence of IdoA 2S→GlcNS. These methods comprise contacting the IdoA 2S→GlcNS-containing polysaccharide preparation with a 3-OST-3 protein in the presence of a sulfate donor under conditions which permit the 3-OST-3 protein to convert the IdoA 2S→GlcNS sequence to IdoA 2S→GlcNS 3S. In particular embodiments, the IdoA 2S→GlcNS sequence comprises a part of a GlcNS→IdoA 2S→GlcNS sequence. The 3-OST proteins for use in these methods

are selected from 3-OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4, ce3-OST, functional fragments of these 3-OSTs having 3-O-sulfotransferase activity, conservative substitution variants of these 3-OSTs having 3-O-sulfotransferase activity, and chimeric 3-OSTs having 3-O-sulfotransferase activity. In preferred embodiments, the sulfate donor is 3'-phospho-adenosine 5'-phosphosulfate (PAPS).

In another aspect, the present invention provides methods for partially sequencing complex polysaccharides such as heparan sulfates or other glycosaminoglycans (GAGs). In these methods, a pool of polysaccharides which includes sequences which may be 3-O-sulfated is contacted with a 3-OST protein in the presence of a sulfate donor (e.g., PAPS) under conditions which permit sulfation by the 3-OST. The treated polysaccharides are then subjected to degradation by enzymes which degrade polysaccharides in a sequence-specific manner (e.g., polysaccharide lyases; heparinase I, II or III; heparitinase) and the size profile of the resulting fragments is determined. An identical pool which has not been treated with 3-OST is similarly cleaved by the same enzymes and a size profile determined. Changes in the size profiles indicate that 3-OST activity has modified the saccharide units so as to prevent (or permit) cleavage at sites which previously were (or were not) cleaved. Thus, comparison of the profiles will indicate positions at which the target sequences for 3-OST activity are present and provide a partial polysaccharide sequence.

In another embodiment, the sequence of complex polysaccharides such as HS or GAGs may be partially determined using sequence specific polysaccharide affinity fractionation. To this end, 3-OST proteins which lack enzymatic function but retain sequence-specific HS or GAG binding capacity can be identified or produced (e.g., altering or deleting a portion of the catalytic ST domain by site-directed mutagenesis or deletion mutagenesis). These inactive forms will bind HS or GAGs in a sequence dependent manner and allow sequence-specific saccharide affinity fractionation from complex mixtures of GAGs. The purified structures may be degraded in a step-wise fashion with exolytic, endolytic enzymes and/or nitrous acid, and the resulting degradation products can be compared to standard compounds of known structure. This method will allow the quantitation and characterization of known structures contained within unknown complex polysaccharide samples.

In another embodiment, partial sequence information can be obtained using the 3-OSTs of the invention or other heparan sulfate sequence specific binding ligands as protective groups prior to treating the HS or GAG with modifying agents that detectably alter the HS or GAG. Useful protective groups include catalytically inactive enzymes, chimeric enzymes and small molecule ligands with identified sequence binding specificities. The protecting group is contacted with the heparan or other glycosaminoglycans (GAGs), and the resultant complex is treated with one or more modifying agents. Useful modifying agents include catalytically active heparan lyases, sulfotransferases, N-deacetylases, N-acetyltransferases, epimerases, or chimeric proteins of the invention. In embodiments where multiple protecting groups and/or modifying reagents are used in combination, the sample is first contacted with the protective group, then one or more modifying reagents may be with contacted with the protected polysaccharide, either simultaneously or in turn. The protective group(s) will interfere with the ability of a modifying agent to interact with, attach to and/or cleave specific GAG sequence motifs. The sample can then be analyzed for ligand-specific protection and/or cleavage to elucidate the sequence of the original GAG using separation and/or quantitation using methods known in the art.

In another aspect, the present invention also provides methods for diagnosing individuals with disorders involving heparan sulfate biosynthesis comprising assaying such individuals for the presence of mutations in 3-OST genes/proteins. Such assays include nucleic acid based assays (employing the nucleic acids of the present invention), protein based assays (employing the antibodies of the present invention), and HS based assays employing the glycosaminoglycan sequencing methods of the present invention.

These and other aspects of the present invention will be apparent to one of ordinary skill in the art from the following detailed description.

#### Brief Description of the Drawings

Fig. 1 is an alignment of the amino acid sequences of murine and human 3-OST-1 proteins showing the high degree of homology. Vertical bars (|) between residues indicate identical residues.

Fig. 2 is an alignment of the sulfotransferase domains of human NST-1, human NST-2, *C. elegans* 3-OST, human 3-OST-4, human 3-OST-3A, human 3-OST-2, and human 3-OST-1.

Fig. 3 is a schematic depiction of the structures of the 3-OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B and 3-OST-4 proteins.

### Detailed Description of the Invention

#### Definitions

In order to more clearly and distinctly point out and describe the subject matter that applicants regard as the invention, the following definitions are provided for certain terms used in the following written description and the appended claims.

Isolated nucleic acids. As used herein with respect to nucleic acids derived from naturally-occurring sequences, the term "isolated nucleic acid" means a ribonucleic or deoxyribonucleic acid which comprises a naturally-occurring nucleotide sequence and which is manipulable by standard recombinant DNA techniques, but which is not covalently joined to the nucleotide sequences that are immediately contiguous on its 5' and 3' ends in the naturally-occurring genome of the organism from which it is derived. As used herein with respect to synthetic nucleic acids, the term "isolated nucleic acid" means a ribonucleic or deoxyribonucleic acid which comprises a nucleotide sequence which does not occur in nature and which is manipulable by standard recombinant DNA techniques. An isolated nucleic acid is manipulable by standard recombinant DNA techniques when it may be used in, for example, amplification by polymerase chain reaction (PCR), *in vitro* translation, ligation to other nucleic acids (e.g., cloning or expression vectors), restriction from other nucleic acids (e.g., cloning or expression vectors), transformation of cells, hybridization screening assays, or the like. The term "isolated nucleic acids" is also intended to embrace synthetic oligonucleotides such as peptide nucleic acids (PNAs), nucleotides joined by phosphorothioate or other non-phosphodiester linkages, nucleic acids incorporating functionally equivalent nucleotide analogs, and the like.

Transformation As used herein, means any method of introducing exogenous a nucleic acid into a cell including, but not limited to, transformation, transfection, electroporation, microinjection, direct injection of naked nucleic acid, particle-mediated delivery, viral-mediated transduction or any other means of delivering a



nucleic acid into a host cell which results in transient or stable expression of said nucleic acid or integration of said nucleic acid into the genome of said host cell or descendant thereof.

Substantially pure. As used herein with respect to protein preparations, the term "substantially pure" means a preparation which contains at least 60% (by dry weight) the protein of interest, exclusive of the weight of other intentionally included compounds. Preferably the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by dry weight the protein of interest, exclusive of the weight of other intentionally included compounds. Purity can be measured by any appropriate method, e.g., column chromatography, gel electrophoresis, or HPLC analysis. If a preparation intentionally includes two or more different proteins of the invention, a "substantially pure" preparation means a preparation in which the total dry weight of the proteins of the invention is at least 60% of the total dry weight, exclusive of the weight of other intentionally included compounds. Preferably, for such preparations containing two or more proteins of the invention, the total weight of the proteins of the invention be at least 75%, more preferably at least 90%, and most preferably at least 99%, of the total dry weight of the preparation, exclusive of the weight of other intentionally included compounds. Thus, if the proteins of the invention are mixed with one or more other proteins (e.g., serum albumin, 6-OST) or compounds (e.g., diluents, detergents, excipients, salts, polysaccharides, sugars, lipids) for purposes of administration, stability, storage, and the like, the weight of such other proteins or compounds is ignored in the calculation of the purity of the preparation.

Similarity. As used herein with respect to amino acid sequences, the "similarity" between two sequences means the percentage of amino acid residue positions, after aligning the sequences according to standard techniques, at which the two sequences have identical or similar residues. In general, "similar" residues include those which are regarded in the art as "conservative substitutions" (see, e.g., Dayhoff et al. (1978), Atlas of Protein Sequence and Structure Vol. 5 (Suppl. 3), pp. 354-352, Natl. Biomed. Res. Found., Washington, D.C.); which fall within the groups (a) methionine, leucine, isoleucine and valine, (b) phenylalanine, tyrosine and tryptophan, (c) lysine, arginine and histidine, (d) alanine and glycine, (e) serine and

threonine, (f) glutamine and asparagine, and (g) glutamate and aspartate; or which are otherwise shown to have no substantial effect on the biological activity of the protein. Numerical values for similarity were determined using the PileUp program. This program performed multiple sequence alignments based on methods of Feng and Doolittle (1987) *J. Mol. Evol.* 35: 351-360, and Higgins and Sharp (1998), *CABIOS* 5:151-153. Using these methods for each sequence alignment, the gap weight was set at 3.0 and the gap length was set at 0.10. Percentages of similarity recited in the appended claims may be determined by these methods.

Chimeric protein. As used herein, the term "chimeric protein" means a protein having an amino acid sequence which is a positionally conserved combination of the amino acid sequences of two or more other proteins. Thus, for a chimera of two or more reference proteins, the amino acid sequences of the reference proteins are aligned by standard techniques to identify residues which correspond at each position, allowing for relative insertions/deletions as necessary. Then, for each amino acid position of the chimeric protein, an amino acid residue is selected from the residues present at corresponding positions in the two or more reference proteins (allowing for no residue in the chimera when deletions are present amongst the reference proteins). The resultant chimera has an amino acid sequence which is a combination of the reference amino acid sequences, in which the relative position of each residue selected from the reference sequences is conserved within the chimera.

Heparan sulfate. As used herein, the term "heparan sulfate" or the abbreviation "HS" means a polysaccharide of the form  $([\rightarrow 4\text{-D-GlcAp}\beta 1 \text{ or } \rightarrow 4\text{-L-IdoAp}\alpha 1] \rightarrow 4\text{-D-GlcNp[Ac or S]}\alpha 1 \rightarrow)_n$  which is modified to a variable extent by sulfation of the 2-*O*-position of Glc and Ido residues, and the 6-*O*- and 3-*O*- positions of GlcN[Ac or S] residues. Therefore, this definition encompasses all glycosaminoglycan compounds referred to as heparan(s), heparan sulfate(s), heparin(s), heparin sulfate(s), heparitin(s), heparitin sulfate(s), heparanoid(s), heparosan(s). The heparan molecules may be pure glycosaminoglycans or can be linked to other molecules including other polymers such as proteins, and lipids, or small molecules such as biotin.

The Heparan Sulfate D-Glucosaminyl 3-O-Sulfotransferases The present invention depends, in part, upon the identification and molecular cloning of cDNAs encoding mammalian heparan sulfate D-glucosaminyl 3-O-sulfotransferases (3-OSTs). These proteins have been designated 3-OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B, and 3-OST-

4. In addition, a nematode 3-OST from *C. elegans*, ce3-OST, has been identified.

3-OST-1s. Disclosed herein are the isolation and identification of murine and human 3-OST-1 cDNAs (SEQ ID NO: 1 and SEQ ID NO: 3, respectively). The coding regions of these cDNAs extend from, respectively, nucleotide positions 323-1255 of SEQ ID NO: 1 and positions 119-1039 of SEQ ID NO : 3. The protein

coding portions of the cDNAs are 85% identical and encode proteins of 311 and 307 amino acids (SEQ ID NO: 2 and SEQ ID NO: 4, respectively) which are 93% similar.

The murine and human protein sequences are aligned in Figure 1. Each protein includes a twenty residue presumptive signal peptide (residues 1-20 of SEQ ID NO: 2 and SEQ ID NO: 4) which is cleaved off to form the mature form of these proteins.

The mouse 3-OST-1 contains an extra four residues (Ala<sup>24</sup>-Pro<sup>25</sup>-Gly<sup>26</sup>-Pro<sup>27</sup>) not found in the human form. Each protein has five potential *N*-glycosylation sites (at residues 52-54, 141-143, 196-198, 246-248 and 253-255 of SEQ ID NO: 2, and residues 48-50, 137-139, 192-194, 242-244, 249-251 of SEQ ID NO: 4). *N*-glycosylation of at least some of these sites appears important to 3-OST protein stability, specificity and/or activity. After the 3-OST-1 signal peptide, there is a domain rich in the residues S, P, L, A, and G (SPLAG-rich domain) (residues 21-52 of SEQ ID NO: 2 and residues 21-48 of SEQ ID NO: 4). 3-OST-1 and all known NST species possess a homologous carboxy terminal sulfotransferase (ST) domain of ~260 amino acids (residues 53-311 of SEQ ID NO: 2 and residues 49-307 of SEQ ID NO:

4) that exhibits homology to all known sulfotransferases and which includes the minimal fragment necessary for sulfation activity. Figure 2 shows a sequence alignment of the ST domains of the sulfotransferases NST-1 (SEQ ID NO: 13), NST-2 (SEQ ID NO: 14), OST-1, OST-2, OST-3A/B, and OST-4. Within this region is a conserved sequence (at residues 260-269 of SEQ ID NO: 2, and 256-265 of SEQ ID NO: 4) which is a presumptive cysteine-bridged peptide loop thought to be involved in heparan sulfate substrate specificity. This cysteine-bridged peptide loop is part of the larger HS-binding domain (residues 250-276 of SEQ ID NO: 2 and 246-272 of

SEQ ID NO: 4). A conserved lysine residue (residue 68 of SEQ ID NO: 2, and 64 of SEQ ID NO: 4) is presumptively catalytic.

The 3-OST-1 proteins have 3-O-sulfotransferase activity on polysaccharide sequences including the sequence GlcA→GlcNS ±6S, and convert this polysaccharide sequence to the sequence to GlcA→GlcNS 3S ±6S. Of particular importance, the 3-OST-1 proteins are useful in converting HS<sup>act</sup> precursor sequences (i.e., IdoA→GlcNAc 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S; or IdoA→GlcNS 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S) to HS<sup>act</sup>. The 3-OST-1 proteins are highly expressed in endothelial cells, brain and kidney tissues, and to a lesser extent in heart, lung, skeletal muscle and placenta. The human 3-OST-1 gene has been syntactically localized to chromosome 4, and more particularly to chromosome segment 4p15-16.

3-OST-2s. Also disclosed herein are the isolation and identification of a human 3-OST-2 cDNA (SEQ ID NO: 5). The coding region of this cDNA extends from nucleotide positions 73-1173 of SEQ ID NO: 5. The cDNA encodes a protein of 367 amino acids (SEQ ID NO: 6). The protein has four potential *N*-glycosylation sites (at residues 102-104, 193-195, 235-237 and 306-308 of SEQ ID NO: 6). *N*-glycosylation of at least some of these sites appears important to 3-OST protein stability, specificity and/or activity. The 3-OST-2 protein has a putative N-terminal cytoplasmic domain (residues 1-19 of SEQ ID NO: 6), followed by a putative transmembrane domain (residues 20-41 of SEQ ID NO: 6), followed by a SPLAG-rich domain (residues 42-109 of SEQ ID NO: 6). This is followed by the characteristic carboxy terminal ST domain of ~260 amino acids (residues 110-367 of SEQ ID NO: 6) that exhibits homology to all known sulfotransferases and which includes the minimal fragment necessary for sulfation activity. Within this region is a conserved sequence (at residues 313-325 of SEQ ID NO: 6) which is a presumptive cysteine-bridged peptide loop thought to be involved in heparan sulfate substrate specificity. This cysteine-bridged peptide loop is part of the larger HS-binding domain (residues 303-332 of SEQ ID NO: 6). A conserved lysine residue (residue 24 of SEQ ID NO: 6) is presumptively catalytic. A cDNA of an allelic variant has also been identified, which includes four silent nucleotide substitutions (G→A at bp 804,

T→G at bp 1249, T→C at bp 1350, and C→T at bp 1507 of SEQ ID NO: 5) which do not affect the encoded protein.

The 3-OST-2 proteins have 3-O-sulfotransferase activity on polysaccharide sequences including the sequences GlcA 2S→GlcNS or GlcNS→GlcA 2S→GlcNS, and convert these polysaccharide sequences to GlcA 2S→GlcNS 3S or GlcNS→GlcA 2S→GlcNS 3S, respectively. The 3-OST-2 proteins are not expressed in endothelial cells, but are highly expressed in brain tissues, and to a lesser extent in heart, lung, skeletal muscle and placenta. The human 3-OST-2 gene has been localized to chromosome 16, and more particularly to chromosome segment 16p12.3.

3-OST-3As. Also disclosed herein are the isolation and identification of a human 3-OST-3A cDNA (SEQ ID NO: 7). The coding region of this cDNA extends from nucleotide positions 799-2016 of SEQ ID NO: 7. The cDNA encodes a protein of 406 amino acids (SEQ ID NO: 8). The protein has two potential *N*-glycosylation sites (at residues 273-275 and 344-346 of SEQ ID NO: 8). *N*-glycosylation of one or more of these sites appears important to 3-OST protein stability, specificity and/or activity. The 3-OST-3A protein has a putative N-terminal cytoplasmic domain (residues 1-24 of SEQ ID NO: 8), followed by a putative transmembrane domain (residues 25-43 of SEQ ID NO: 8), followed by a SPLAG-rich domain (residues 44-147 of SEQ ID NO: 8). This is followed by the characteristic carboxy terminal ST domain of ~260 amino acids (residues 148-406 of SEQ ID NO: 8) that exhibits homology to all known sulfotransferases and which includes the minimal fragment necessary for sulfation activity. Within this region is a conserved sequence (at residues 351-363 of SEQ ID NO: 8) which is a presumptive cysteine-bridged peptide loop thought to be involved in heparan sulfate substrate specificity. This cysteine-bridged peptide loop is part of the larger HS-binding domain (residues 341-370 of SEQ ID NO: 8). A conserved lysine residue (residue 162 of SEQ ID NO: 8) is presumptively catalytic.

The 3-OST-3A proteins have 3-O-sulfotransferase activity on polysaccharide sequences including the sequences IdoA 2S→GlcNS or GlcNS→IdoA 2S→GlcNS, and convert these polysaccharide sequences to IdoA 2S→GlcNS 3S or GlcNS→IdoA 2S→GlcNS 3S, respectively. The 3-OST-3A proteins are not expressed in endothelial

cells, but are highly expressed in kidney, placenta and liver tissues, and to a lesser extent in brain, heart, lung, and skeletal muscle.

3-OST-3Bs. Also disclosed herein are the isolation and identification of a human 3-OST-3B cDNA (SEQ ID NO: 9). The coding region of this cDNA extends from nucleotide positions 331-1500 of SEQ ID NO: 9. The cDNA encodes a protein of 390 amino acids (SEQ ID NO: 10). The protein has two potential *N*-glycosylation sites (at residues 258-260 and 329-331 of SEQ ID NO: 10). *N*-glycosylation of one or more of these sites appears important to 3-OST protein stability, specificity and/or activity. The 3-OST-3B protein has a putative N-terminal cytoplasmic domain (residues 1-32 of SEQ ID NO: 10), followed by a putative transmembrane domain (residues 33-65 of SEQ ID NO: 10), followed by a SPLAG-rich domain (residues 66-132 of SEQ ID NO: 10). This is followed by the characteristic carboxy terminal ST domain of ~260 amino acids (residues 133-390 of SEQ ID NO: 10) that exhibits homology to all known sulfotransferases and which includes the minimal fragment necessary for sulfation activity. Within this region is a conserved sequence (at residues 336-348 of SEQ ID NO: 10) which is a presumptive cysteine-bridged peptide loop thought to be involved in heparan sulfate substrate specificity. This cysteine-bridged peptide loop is part of the larger HS-binding domain (residues 326-355 of SEQ ID NO: 10). A conserved lysine residue (residue 147 of SEQ ID NO: 10) is presumptively catalytic.

The 3-OST-3B proteins have 3-O-sulfotransferase activity on polysaccharide sequences including the sequences IdoA 2S→GlcNS or GlcNS→IdoA 2S→GlcNS, and convert these polysaccharide sequences to IdoA 2S→GlcNS 3S or GlcNS→IdoA 2S→GlcNS 3S, respectively. The 3-OST-3A proteins are not expressed in endothelial cells, but are highly expressed in kidney, placenta and liver tissues, and to a lesser extent in brain, heart, lung, and skeletal muscle.

3-OST-4s. Also disclosed herein are the isolation and identification of a human 3-OST-4 nucleic acid sequence (SEQ ID NO: 11). This sequence represents is a possible or predicted heteronuclear RNA species, and is a composite of 5' genomic sequences information and an overlapping partial cDNA. The coding region of this sequence extends from nucleotide positions 847-2214 of SEQ ID NO: 11, and encodes a protein of 456 amino acids (SEQ ID NO: 12). The protein has two

potential *N*-glycosylation sites (at residues 318-320 and 389-391 of SEQ ID NO: 12). *N*-glycosylation of one or more of these sites appears important to 3-OST protein stability, specificity and/or activity. The 3-OST-4 includes the characteristic carboxy terminal ST domain of ~260 residues (residues 207-456 of SEQ ID NO: 12) that exhibits homology to all known sulfotransferases and which includes the minimal fragment necessary for sulfation activity. Within this region is a conserved sequence (at residues 396-408 of SEQ ID NO: 12) which is a presumptive cysteine-bridged peptide loop thought to be positioned near the active site. This cysteine-bridged peptide loop is part of the larger HS-binding domain (residues 386-415 of SEQ ID NO: 12). A conserved lysine residue (residue 207 of SEQ ID NO: 12) is presumptively catalytic.

The 3-OST-4 proteins have sulfotransferase activity, but the sequence specificity of this activity has not yet been determined. The 3-OST-4 proteins appear to be expressed at detectable levels only in the brain. The human 3-OST-4 gene has been localized to chromosome 16, and more particularly to chromosome segment 16p11.

*C. elegans* 3-OSTs. Also disclosed herein is the identification of a *C. elegans* homologue of the human 3-OSTs, ce3-OST. This protein is disclosed as SEQ ID NO: 15, and includes the characteristic carboxy terminal ST domain of ~260 residues (residues 23-291 of SEQ ID NO: 15) that exhibits homology to all known sulfotransferases and which includes the minimal fragment necessary for sulfation activity. Within this region is a conserved sequence (at residues 240-250 of SEQ ID NO: 15) which is a presumptive cysteine-bridged peptide loop thought to be positioned near the active site. This cysteine-bridged peptide loop is part of the larger HS-binding domain (residues 230-257 of SEQ ID NO: 15). A conserved lysine residue (residue 38 of SEQ ID NO: 15) is presumptively catalytic.

The *C. elegans* 3-OST proteins have sulfotransferase activity, but the sequence specificity of this activity has not yet been determined. BLAST and Genefinder analysis of genomic cosmids predicts that ce3-OST is an intraluminal resident protein of 291 residues encoded by 4 exons (clone F52B10, Gban U41990; residues 26317-26090, 21886-21732, 21682-21395, and 21345-21140).

The homology between the sulfotransferase domain of the ce3-OST and the human 3-OST and NST proteins is illustrated in Fig. 2. Based on this sequence alignment, one may also produce chimeric proteins between and the *C elegans* protein and its human homologues.

5 Isolated Nucleic Acids

In one aspect, the present invention provides isolated nucleic acids encoding 3-OST proteins or functional fragments thereof. In preferred embodiments, the 3-OST proteins are 3-OST-1 proteins, 3-OST-2 proteins, 3-OST-3A proteins, 3-OST-3B proteins, 3-OST-4 proteins, or ce3-OST proteins. In particularly preferred  
10 embodiments, the 3-OST proteins are those disclosed as SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 15. As shown in the examples below, the isolated nucleic acids encoding all or a portion of one mammalian 3-OST protein may be used to isolate homologues in other species by standard techniques known to those of ordinary skill in the art. Thus, the  
15 present invention also enables isolated nucleic acids encoding the 3-OST proteins of other mammalian species including, for example, rats, goats, sheep, cows, pigs, and non-human primates. Similarly, the isolated nucleic acids disclosed herein may be used to screen additional human or other mammalian genetic libraries (e.g., genomic or cDNA libraries) to identify allelic variants of the particularly disclosed sequences.  
20 Thus, the present invention also enables isolated nucleic acids encoding human and other mammalian 3-OST allelic variants.

In another aspect, the present invention provides isolated nucleic acids encoding functional fragments of 3-OST proteins, 3-OST protein variants in which conservative substitutions have been made for certain residues, or encoding chimeric  
25 3-OST proteins in which the sequences of two or more 3-OST proteins have been mixed, to produce non-naturally occurring variants which retain sequence-specific HS binding affinity and/or 3-O-sulfotransferase activity. The preferred amino acid sequences of such variants are described below.

In preferred embodiments, the isolated nucleic acids encoding a mammalian 3-OST or functional fragment thereof have at least 60%, preferably at least 70%, and  
30 more preferably at least 80% nucleotide sequence identity to the coding regions of the mammalian 3-OST sequences particularly disclosed herein (SEQ ID NO: 1, SEQ ID



NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9 and SEQ ID NO: 11), and encode at least a functional fragment having sequence-specific HS binding affinity and/or 3-O-sulfotransferase activity. Most preferably, the sequences have at least 90% or 95% nucleotide sequence identity to the disclosed reference sequences.

5 As will be apparent to one of ordinary skill in the art, the degeneracy of the genetic code allows for numerous nucleotide substitutions in a given coding sequence which do not affect the amino acid sequence of the encoded protein. Thus, the present invention also provides for isolated nucleic acids which differ from any of the above-described sequences only by the substitution of such synonymous codons.

10 The isolated nucleic acids of the present invention may be joined to other nucleic acid sequences for use in various applications. Thus, for example, the isolated nucleic acids of the invention may be ligated into cloning or expression vectors, as are commonly known in the art and as described in the examples below. In addition, the nucleic acids of the invention may be joined in-frame to sequences encoding another  
15 polypeptide so as to form a fusion protein, as is commonly known in the art and as described in the examples below. Thus, in certain embodiments, the present invention provides cloning, expression and fusion vectors comprising any of the above-described nucleic acids.

In another aspect, the isolated nucleic acids of the present invention may  
20 comprise only a portion of a nucleotide sequence encoding a complete mammalian 3-OST protein. For example, and as described more fully below, the 3-OST-1 proteins comprise a signal sequence which is removed post-translationally to yield the mature proteins. In some instances (e.g., when translating 3-OST-1 proteins *in vitro*), it may be preferable to employ an isolated nucleic acid which encodes only the mature  
25 protein. In addition, the four C-terminal residues of 3-OST-1 are believed to be involved in localization of the protein within the Golgi apparatus. In some instances (e.g., when encoding 3-OST-1 proteins for use *in vitro*), it may be preferable to employ an isolated nucleic acid which does not encode these residues, as they will be unnecessary for *in vitro* function. As described above, an approximately 260 residue  
30 portion of the 3-OST proteins includes the catalytically active region (ST domain) and, therefore, it may be preferable to employ an isolated nucleic acid which encodes only this functional fragment which retains 3-O-sulfotransferase activity. Thus, in

certain preferred embodiments, the present invention provides isolated nucleic acid sequences encoding mature forms of a mammalian 3-OST-1 protein, C-terminally truncated forms of the 3-OST proteins, or minimal functional fragments of the 3-OST proteins. In addition, as described above, these sequences may also encode  
5 conservative substitution variants or chimeras of 3-OST proteins, and may include synonymous codon substitutions.

In another aspect, the present invention provides for nucleic acids which comprise a sequence of at least 16-18, preferably 18-20 consecutive nucleotides from any one of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID  
10 NO: 9 and SEQ ID NO: 11. Such nucleic acid sequences have utility for determining the levels of expression of 3-OST transcripts in cells or tissues, for identifying tissues in which the 3-OST genes are differentially expressed (see above), for encoding peptide fragments which may be used to raise antibodies to corresponding regions of the 3-OST proteins, identifying chromosomes bearing the corresponding 3-OST  
15 sequences (see above), for priming polymerase chain reaction amplification of 3-OST sequences (e.g., prior to *in vitro* translation, see below), and for various other utilities which will be apparent to those skilled in the art. Particularly preferred sequences for PCR amplification include those which are 5' to and/or include the initiation codon, which are 5' to and/or include the codons encoding the signal peptide cleavage site, or  
20 which are 3' to and/or include the termination codon. Sequences useful for encoding peptide fragments include those which are located within the coding region.

#### Cell Lines and Transgenic Animals

The present invention also provides for cells or cell lines, both prokaryotic and eukaryotic, into which have been introduced the nucleic acids of the present invention  
25 so as to cause clonal propagation of those nucleic acids and/or expression of the proteins or peptides encoded thereby. Such cells or cell lines have utility in the propagation and production of the nucleic acids of the invention, as well as the production of the proteins of the present invention. As used herein, the term "transformed cell" is intended to embrace any cell, or the descendant of any cell, into  
30 which has been introduced any of the nucleic acids of the invention, whether by transformation, transfection, transduction, infection, or other means. Methods of producing appropriate vectors, transforming cells with those vectors, and identifying

transformants are well known in the art and are only briefly reviewed here (see, for example, Sambrook et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).

Prokaryotic cells useful for producing the transformed cells of the invention include members of the bacterial genera *Escherichia* (e.g., *E. coli*), *Pseudomonas* (e.g., *P. aeruginosa*), and *Bacillus* (e.g., *B. subtilis*, *B. stearothermophilus*), as well as many others well known and frequently used in the art. Prokaryotic cells are particularly useful for the production of large quantities of the proteins or peptides of the invention (e.g., naturally occurring or synthetic 3-OSTs, fragments of the 3-OSTs, fusion proteins of the 3-OSTs). Bacterial cells (e.g., *E. coli*) may be used with a variety of expression vector systems including, for example, plasmids with the T7 RNA polymerase/promoter system, bacteriophage  $\lambda$  regulatory sequences, or M13 Phage regulatory elements. Bacterial hosts may also be transformed with fusion protein vectors which create, for example, Protein A, lacZ, trpE, maltose-binding protein, poly-His tag, or glutathione-S-transferase fusion proteins. All of these, as well as many other prokaryotic expression systems, are well known in the art and widely available commercially (e.g., pGEX-27 (Amrad, USA) for GST fusions).

Eukaryotic cells and cell lines useful for producing the transformed cells of the invention include mammalian cells (e.g., endothelial cells, mast cells, COS cells, CHO cells, fibroblasts, hybridomas, oocytes, embryonic stem cells), insect cells lines (e.g., *Drosophila* Schneider cells), yeast, and fungi. Eukaryotic cells are particularly useful for embodiments in which it is necessary that the 3-OST proteins, or functional fragments thereof, be properly post-translationally modified (e.g., *N*-glycosylated) because *N*-glycosylation of these proteins appears to be important to their stability and/or activity. Currently preferred cells are mammalian cells and, in particular, COS-7 cells, CHO cells, murine primary cardiac microvascular endothelial cells (CME), murine mast cell line C57.1, human primary endothelial cells of umbilical vein (HUVEC), F9 embryonal carcinoma cells, rat fat pad endothelial cells (RFPEC), L cells (e.g., murine LTA *tk*<sup>-</sup> cells), and cells derived from the transgenic animals of the invention.

To accomplish expression in eukaryotic cells, a wide variety of vectors have been developed and are commercially available which allow inducible (e.g.,

LacSwitch expression vectors, Stratagene, La Jolla, CA) or constitutive (e.g., pcDNA3 vectors, Invitrogen, Chatsworth, CA) expression of 3-OST nucleotide sequences under the regulation of an artificial promoter element. Such promoter elements are often derived from CMV or SV40 viral genes, although other strong promoter elements which are active in eukaryotic cells can also be employed to induce transcription of 3-OST nucleotide sequences. Typically, these vectors also contain an artificial polyadenylation sequence and 3' UTR which can also be derived from exogenous viral gene sequences or from other eukaryotic genes. These expression systems are commonly available from commercial sources and are typified by vectors such as pcDNA3 and pZeoSV (Invitrogen, San Diego, CA). As described below, the vector pcDNA3 has been successfully used to cause expression of 3-OST-1 proteins in transfected COS-7 cells. Numerous expression vectors are available from commercial sources to allow expression of any desired 3-OST transcript in more or less any desired cell type, either constitutively or after exposure to a certain exogenous stimulus (e.g., withdrawal of tetracycline or exposure to IPTG).

Vectors may be introduced into the recipient or "host" cells by various methods well known in the art including, but not limited to, calcium phosphate transfection, strontium phosphate transfection, DEAE dextran transfection, electroporation, lipofection, microinjection, ballistic insertion on micro-beads, protoplast fusion or, for viral or phage vectors, by infection with the recombinant virus or phage.

#### Transgenic Animal Models

The present invention also provides for the production of transgenic non-human animal models in which wild type, allelic variant, chimeric, or antisense 3-OST sequences are expressed, or in which 3-OST sequences have been inactivated or deleted (e.g., "knock-out" constructs) or replaced with reporter or marker genes (e.g., "knock-in reporter construct"). The 3-OST sequences may be conspecific to the transgenic animal (e.g., murine sequences in a transgenic mouse) or transspecific to the transgenic animal (e.g. human sequence in a transgenic mouse). In such a transgenic animal, the transgenic sequences may be expressed inducibly, constitutively or ectopically. Expression may be tissue-specific or organism-wide. Engineered expression of 3-OST sequences in tissues and cells not normally containing 3-OST

gene products may cause novel alterations of heparan polysaccharide structure and lead to novel cell or tissue phenotypes. Ectopic or altered levels of expression of 3-OST sequences may alter cell, tissue and/or developmental phenotypes. Transgenic animals are useful as models of thromboembolic and other disorders arising from defects in heparan sulfate biosynthesis or metabolism. Transgenic animals are also useful for screening compounds for their effects on HS biosynthesis mediated by 3-OSTs. Transgenic animals transformed with reporter constructs may be used to measure the transcriptional effects of small molecules, drugs, protein physiological mediators, carbohydrate effectors, mimetic compounds or physical perturbations on the expression of 3-OST loci *in vivo*. The transgenic animals of the invention, may be used to screen such compounds for therapeutic utility.

Animal species suitable for use in the animal models of the present invention include, but are not limited to, rats, mice, hamsters, guinea pigs, rabbits, dogs, cats, goats, sheep, pigs, and non-human primates (e.g., Rhesus monkeys, chimpanzees). For initial studies, transgenic rodents (e.g., mice) are preferred due to their relative ease of maintenance and shorter life spans. Transgenic non-human primates may be preferred for longer term studies due to their greater similarity to humans and their higher cognitive abilities.

Using the a nucleic acid disclosed and otherwise enabled herein, there are now several available approaches for the creation of a transgenic animal. Thus, the enabled animal models include: (1) animals in which sequences encoding at least a functional fragment of a wild type 3-OST gene has been recombinantly introduced into the genome of the animal as an additional gene, under the regulation of either an exogenous or an endogenous promoter element, and as either a minigene (i.e., a genetic construct of the 3-OST with the introns, if any, removed) or a large genomic fragment; (2) animals in which sequences encoding at least a functional fragment of a normal 3-OST gene have been recombinantly substituted for one or both copies of the animal's homologous 3-OST gene by homologous recombination or gene targeting; (3) animals in which one or both copies of one of the animal's homologous 3-OST genes have been recombinantly "humanized" by the partial substitution of sequences encoding the human homologue by homologous recombination or gene targeting; (4) animals in which sequences encoding 3-OST transcriptional elements linked to a

reporter gene have replaced the endogenous 3-OST gene and transcriptional elements;  
(5) "knock-out" animals in which one or both copies of the animal's 3-OST sequences  
have been partially or completely deleted or have been inactivated by the insertion or  
substitution by homologous recombination or gene targeting of exogenous sequences  
(e.g., stop codons); (6) animals in which additional genes related to the biosynthesis  
or metabolism of heparan sulfates have been altered (e.g., a murine transgenic in  
which all of the genes in the HS pathway have been humanized). These and other  
transgenic animals of the invention are useful as models of thromboembolic and other  
disorders arising from defects in heparan sulfate biosynthesis or metabolism. These  
animals are also useful for screening compounds for their effects on HS biosynthesis  
mediated by 3-OSTs.

To produce an animal model (e.g., a transgenic mouse), a wild type or  
allelic variant 3-OST sequence or a wild type or allelic variant of a recombinant  
nucleic acid encoding at least a functional fragment of a 3-OST is preferably inserted  
into a germ line or stem cell using standard techniques of oocyte or embryonic stem  
cell microinjection, or other form of transformation of such cells. Alternatively, other  
cells from adult organism may be employed. Animals produced by these or similar  
processes are referred to as transgenic. Similarly, if it is desired to inactivate or  
replace an endogenous 3-OST sequence, homologous recombination using oocytes,  
embryonic stem or other cells may be employed. Animals produced by these or  
similar processes are referred to as "knock-out" (inactivation) or "knock-in"  
(replacement) models.

For oocyte injection, one or more copies of the recombinant DNA  
constructs of the present invention may be inserted into the pronucleus of a just-  
fertilized oocyte. This oocyte is then reimplanted into a pseudo-pregnant foster  
mother. The liveborn animals are screened for integrants using analysis of DNA (e.g.,  
from the tail veins of offspring mice) for the presence of the inserted recombinant  
transgene sequences. The transgene may be either a complete genomic sequence  
introduced into a host as a YAC, BAC or other chromosome DNA fragment, a cDNA  
with either the natural promoter or a heterologous promoter, or a minigene containing  
all of the coding region and other elements found to be necessary for optimum  
expression.

To create a transgene, the target sequence of interest (e.g., wild type or allelic variant 3-OST sequences) are typically ligated into a cloning site located downstream of some promoter element which will regulate the expression of RNA from the sequence. Downstream of the coding sequence, there is typically an artificial polyadenylation sequence. An alternative approach to creating a transgene is to use an exogenous promoter and regulatory sequences to drive expression of the transgene. Finally, it is possible to create transgenes using large genomic DNA fragments such as YACs which contain the entire desired gene as well as its appropriate regulatory sequences.

Animal models may be created by targeting endogenous 3-OST sequence in order to alter the endogenous sequence by homologous recombination. These targeting events can have the effect of removing endogenous sequence (knock-out) or altering the endogenous sequence to create an amino acid change associated with human disease or an otherwise abnormal sequence (e.g., a sequence which is more like the human sequence than the original animal sequence) (knock-in animal models). A large number of vectors are available to accomplish this and appropriate sources of genomic DNA for mouse and other animal genomes to be targeted are commercially available from companies such as GenomeSystems Inc. (St. Louis, Missouri, USA). The typical feature of these targeting vector constructs is that 2 to 4 kb of genomic DNA is ligated 5' to a selectable marker (e.g., a bacterial neomycin resistance gene under its own promoter element termed a "neomycin cassette"). A second DNA fragment from the gene of interest is then ligated downstream of the neomycin cassette but upstream of a second selectable marker (e.g., thymidine kinase). The DNA fragments are chosen such that mutant sequences can be introduced into the germ line of the targeted animal by homologous replacement of the endogenous sequences by either one of the sequences included in the vector. Alternatively, the sequences can be chosen to cause deletion of sequences that would normally reside between the left and right arms of the vector surrounding the neomycin cassette. The former is known as a knock-in, the latter is known as a knock-out.

Retroviral infection of early embryos can also be done to insert the recombinant DNA constructs of the invention. In this method, the transgene (e.g., a

wild type or allelic variant 3-OST sequence) is inserted into a retroviral vector which is used to directly infect embryos (e.g., mouse or non-human primate embryos) during the early stages of development to generate partially transgenic animals, some of which bear the transgenes in germline cells.

Alternatively, homologous recombination using a population of stem cells allows for the screening of the population for successful transformants. Once identified, these can be injected into blastocysts, and a proportion of the resulting animals will show germline transmission of the transgene.

Techniques of generating transgenic animals, as well as techniques for homologous recombination or gene targeting, are now widely accepted and practiced. A laboratory manual on the manipulation of the mouse embryo, for example, is available detailing standard laboratory techniques for the production of transgenic mice (69).

Finally, equivalents of transgenic animals, including animals with mutated or inactivated 3-OST sequences may be produced using chemical or x-ray mutagenesis of gametes, followed by fertilization. Using the isolated a nucleic acid disclosed or otherwise enabled herein, one of ordinary skill may more rapidly screen the resulting offspring by, for example, direct sequencing, SSCP, RFLP, PCR, or hybridization analysis to detect mutants, or Southern blotting to demonstrate loss of one allele by dosage.

#### Substantially Pure Proteins

In one aspect, the present invention provides substantially pure preparations of 3-OST proteins. In preferred embodiments, the 3-OST proteins are 3-OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4 or ce3-OST proteins. In particularly preferred embodiments, the 3-OST proteins are those disclosed as SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 15. As shown in the examples below, nucleic acids encoding all or a portion of one mammalian 3-OST protein may be used to isolate homologues in other species by standard techniques known to those of ordinary skill in the art. Thus, the present invention also enables substantially pure protein preparations of 3-OST proteins of other mammalian species including, for example, rats, goats, sheep, cows, pigs, and non-human primates. Similarly, the isolated nucleic acids disclosed herein may be



used to screen additional human or other mammalian genetic libraries (e.g., genomic or cDNA libraries) to identify allelic variants of the particularly disclosed sequences. Thus, the present invention also enables substantially pure protein preparations of human and other mammalian 3-OST allelic variants.

5 In another aspect, the present invention provides 3-OST protein variants in which conservative substitutions have been made for certain residues, or chimeric 3-OST proteins in which the sequences of various 3-OST proteins have been mixed, to produce non-naturally occurring variants which retain 3-O-sulfotransferase activity. Conservative substitutions are preferably made in those regions of the proteins which  
10 are already known to vary amongst the human and murine sequences (see Figure 1) or between the 3-OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B 3-OST-4, and ce3-OST proteins (see, e.g., Figure 2). Substitutions are to be avoided in those areas which have been implicated in catalysis (see above). Chimeric 3-OST proteins may be made using the disclosed sequences as reference sequences, and these chimeras may also be  
15 subjected to conservative substitutions as described above. In addition, based upon the homologies of the 3-OST proteins to other glucosaminyl sulfotransferases (e.g., 2-OST, NST-1, NST-2), one of ordinary skill in the art may produce chimeric 3-OSTs using those proteins as reference sequences (see, e.g., Figure 2).

In preferred embodiments, the 3-OST proteins have at least 60%, , preferably  
20 at least 70%, and more preferably at least 80% amino acid sequence similarity to the mammalian 3-OST sequences particularly disclosed herein, and retain 3-O-sulfotransferase activity. Most preferably, the sequences have at least 90% or 95% amino acid sequence similarity to the disclosed reference sequences. Such sequences may be routinely produced by those of ordinary skill in the art, and 3-O-  
25 sulfotransferase activity may be tested by routine methods such as those disclosed herein.

The substantially pure proteins of the present invention may be joined to other polypeptide sequences for use in various applications. Thus, for example, the proteins of the invention may be joined to one or more additional polypeptides so as to form a  
30 fusion protein, as is commonly known in the art and as described in the examples below. The additional polypeptides may be joined to the N-terminus, C-terminus or both termini of the 3-OST protein. Such fusion proteins may be particularly useful if

the additional polypeptide sequences are easily identified (e.g., by providing an antigenic determinant) or easily purified (e.g., by providing a ligand for affinity purification).

In another aspect, the substantially pure 3-OST proteins of the present invention may comprise only a portion or fragment of the amino acid sequence of a complete mammalian 3-OST protein. For example, as described above, the 3-OST-1 proteins comprise a twenty amino acid signal sequence which is removed post-translationally to yield the mature proteins. In some instances (e.g., when employing 3-OST-1 proteins *in vitro*), it may be preferable to employ only the mature protein or a minimal fragment retaining 3-O-sulfotransferase activity. In addition, the four C-terminal residues of 3-OST-1 may be involved in localization of the protein within the Golgi apparatus. In some instances (e.g., when employing 3-OST-1 proteins *in vitro*), it may be preferable to employ a 3-OST-1 protein which does not include these residues, as they will be unnecessary for *in vitro* function. As described above, an approximately 260 amino acid portion of the 3-OST proteins includes the catalytically active region and, therefore, it may be preferable to employ a 3-OST protein which includes only this functional fragment which retains 3-O-sulfotransferase activity. Thus, in certain preferred embodiments, the present invention provides substantially pure 3-OST proteins including mature forms of a mammalian 3-OST-1 protein, C-terminally truncated forms, or minimal functional fragments thereof. In addition, as described above, these proteins may also comprise conservative substitution variants or chimeras of 3-OST proteins.

In another aspect, the present invention provides for substantially pure protein preparations which comprise a sequence of at least 6-12, preferably 10-16, more preferably 16-22 consecutive amino acid residues from any one of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, and SEQ ID NO: 15. Such polypeptides have utility to raise antibodies to corresponding regions of the 3-OST proteins. In particular, an analysis of the amino acid sequences of the 3-OST proteins suggests that there are regions which will have particular utility in generating antibodies. Thus, in preferred embodiments, the inventions provides antigenic 3-OST polypeptides selected from the group consisting of (a) residues 4-29, 144-152, 208-222, 31-42, 155-181, 72-94, 195-205, 278-293, 113-136, 56-66, 230-

245, 257-263, 301-306, 267-272 and 101-107 of SEQ ID NO: 2; (b) residues 4-22, 140-148, 205-218, 68-90, 191-201, 274-289, 110-133, 51-62, 226-241, 253-259, 151-163, 168-181, 297-302, 27-34, 97-107 and 263-268 of SEQ ID NO: 4; (c) residues 18-44, 199-207, 114-123, 319-328, 250-275, 238-246, 128-143, 47-59, 83-98, 332-349, 178-186, 289-295, 310-316, 63-76, 4-9, 209-218, 170-176 and 300-305 of SEQ ID NO: 6; (d) residues 22-57, 236-256, 166-186, 151-161, 138-147, 77-85, 348-354, 87-94, 323-335, 360-366, 284-314, 217-224, 376-383, 4-20, 130-136, 67-73, 389-395 and 338-343 of SEQ ID NO: 8; (e) residues 221-241, 8-66, 151-171, 135-146, 333-339, 308-320, 345-351, 269-299, 202-209, 361-368, 86-100, 71-80, 115-129, 374-380 and 323-328 of SEQ ID NO: 10; and (f) residues 280-290, 321-364, 371-388, 211-231, 393-399, 310-316, 421-438, 405-411, 262-268 and 292-301 of SEQ ID NO: 12. Note that these polypeptides are listed in decreasing order of preference within in group (a) to (f). Preferred antigenic peptide sequences also include residues 218-231, 87-100, 167-180 and 275-288 of SEQ ID NO: 2, which have been successfully used to generate antibodies to m3-OST-1.

Thus, in another aspect, the present invention provides for antibodies and methods for making antibodies which selectively bind with the 3-OST proteins. These antibodies include monoclonal and polyclonal antibodies, as well as functional antibody fragments such as F(ab) and Fc.

The proteins or peptides of the invention may be substantially purified by any of a variety of methods selected on the basis of the properties revealed by their protein sequences. As shown in the examples below, and previously described (26), cells naturally expressing 3-OST-1 proteins secrete the protein when grown in culture, and the proteins may be isolated from the cell culture medium. The 3-OST-2, 3-OST-3A, 3-OST-3B and 3-OST-4 proteins, however, appear to include transmembrane domains. Thus, these proteins are not expected to be secreted at high levels. Because the 3-OSTs are found in the Golgi apparatus and microsomal bodies of cells which naturally express them, a fraction of cells including these organelles may be isolated and the proteins may be extracted from this fraction by, for example, detergent solubilization. Alternatively the 3-OST proteins, fusion proteins, or fragments thereof, may be purified from cells transformed or transfected with expression vectors. For example, insect cells such as *Drosophila* Schneider cells and baculovirus

expression systems may be employed with vectors such as pBLUEBAC and pMELBAC (Stratagene, La Jolla, CA); yeast expression systems with vectors such as pYESHIS Xpress vectors (Invitrogen, San Diego, CA); eukaryotic expression systems with vectors such as pcDNA3 (Invitrogen, San Diego, CA), which causes constitutive expression, or LacSwitch (Stratagene, La Jolla, CA) which is inducible; or prokaryotic expression systems with vectors such as pKK233-3 (Clontech, Palo Alto, CA). In the event that the protein or fragment localizes within microsomes derived from the Golgi apparatus, endoplasmic reticulum, or other membrane containing structures of such cells, the protein may be purified from the appropriate cell fraction. Alternatively, if the protein does not localize within these structures, or aggregates in inclusion bodies within the recombinant cells (e.g., prokaryotic cells), the protein may be purified from whole lysed cells or from solubilized inclusion bodies by standard means.

Purification can be achieved using standard protein purification procedures including, but not limited to, affinity chromatography, gel-filtration chromatography, ion-exchange chromatography, high-performance liquid chromatography (RP-HPLC, ion-exchange HPLC, size-exclusion HPLC), high-performance chromatofocusing chromatography, hydrophobic interaction chromatography, immunoprecipitation, or immunoaffinity purification. Gel electrophoresis (e.g., PAGE, SDS-PAGE) can also be used to isolate a protein or peptide based on its molecular weight, charge properties and hydrophobicity.

A 3-OST protein, or a fragment thereof, may also be conveniently purified by creating a fusion protein including the desired 3-OST sequence fused to another peptide such as an antigenic determinant (e.g., from Protein A, see below) or poly-His tag (e.g., QIAexpress vectors, QIAGEN Corp., Chatsworth, CA), or a larger protein (e.g., GST using the pGEX-27 vector (Amrad, USA) or green fluorescent protein using the Green Lantern vector (GIBCO/BRL, Gaithersburg, MD). The fusion protein may be expressed and recovered from prokaryotic or eukaryotic cells and purified by any standard method based upon the fusion vector sequence. For example, the fusion protein may be purified by immunoaffinity or immunoprecipitation with an antibody to the non-3-OST portion of the fusion or, in the case of a poly-His tag, by affinity binding to a nickel column. The desired 3-OST protein or fragment can then be further purified from the fusion protein by enzymatic cleavage of the fusion protein.

Methods for preparing and using such fusion constructs for the purification of proteins are well known in the art and numerous kits are now commercially available for this purpose.

Currently preferred methods for small scale purification of 3-OST-1 proteins from the media of LTA cells grown in culture may be found in Liu et al. (26), and methods for purification of 3-OSTs produced recombinantly in COS-7 cells, CHO cells, murine primary cardiac microvascular endothelial cells (CME), murine mast cell line C57.1, and human primary endothelial cells of umbilical vein (HUVEC) may be found in the examples below. These methods may also be adapted for use with other cell and expression systems to obtain substantially pure 3-OST proteins.

In another aspect, the present invention provides for methods for producing the above-described proteins. Thus, in one set of embodiments, the isolated nucleic acids of the invention may be used to transform host cells or create transgenic animals. The proteins of the invention may then be substantially purified by well known methods including, but not limited to, those described in the examples below. Alternatively, the isolated nucleic acids of the invention may be used in cell-free *in vitro* translation systems. Such systems are also well known in the art and include, but are not limited to, that described in the examples below.

#### Antibodies

The present invention also provides antibodies and methods of making antibodies, which will selectively bind to and, thereby, isolate or identify wild type and/or variant forms of the 3-OST proteins. The antibodies of the invention have utility as laboratory reagents for, inter alia, immunoaffinity purification of the 3-OSTs, immunoaffinity purification of 3-OST conjugates or complexes (e.g., 3-OST-AT, 3-OST-HS), Western blotting to identify cells or tissues expressing the 3-OSTs, and immunocytochemistry or immunofluorescence techniques to establish the cellular or extracellular location of the protein.

The antibodies of the invention may be generated using the entire 3-OST proteins of the invention or using any 3-OST epitope which is characteristic of that protein and which substantially distinguishes it from other host proteins. Such epitopes may be identified by comparing sequences of amino acid residues from a 3-

OST sequence to computer databases of protein sequences from the relevant host. Preferably, the epitopes are chosen so as to be highly immunogenic and specific.

In a preferred embodiment, the immunogen/epitope is a protein sequence of at least 6-12, preferably 10-16, more preferably 16-22 consecutive amino acid residues of the disclosed OST genes. In particular, an analysis of the amino acid sequences of the 3-OST proteins suggests that there are regions which will have particular utility in generating antibodies. Thus, in preferred embodiments, the inventions provides antigenic 3-OST polypeptides.

3-OST immunogen preparations may be produced from crude extracts (e.g., microsomal fractions of cells expressing the proteins), from proteins or peptides substantially purified from cells which naturally or recombinantly express them or, for small immunogens, by chemical peptide synthesis. The 3-OST immunogens may also be in the form of a fusion protein in which the non-3-OST region is chosen for its adjuvant properties and/or the ability to either and/or facilitate purification. As used herein, a 3-OST immunogen shall be defined as a preparation including a peptide comprising at least 4-8, and preferably at least 9-15 consecutive amino acid residues of the 3-OST proteins or nucleic acids encoding such a peptide coupled with transcriptional elements, as disclosed or otherwise enabled herein. Therefore, any 3-OST derived polypeptide or protein sequences which are employed to generate antibodies to the 3-OSTs should be regarded as 3-OST immunogens.

The antibodies of the invention may be polyclonal or monoclonal, or may be antibody fragments, including Fab fragments,  $F(ab')_2$ , and single chain antibody fragments. In addition, after identifying useful antibodies by the method of the invention, recombinant antibodies may be generated, including any of the antibody fragments listed above, as well as humanized antibodies based upon non-human antibodies to the 3-OST proteins. In light of the present disclosures of 3-OST proteins, as well as the characterization of other 3-OSTs enabled herein, one of ordinary skill in the art may produce the above-described antibodies by any of a variety of standard means well known in the art. For an overview of antibody techniques, see Antibody Engineering, 2nd Ed., Borrebaek, ed., Oxford University Press, Oxford (1995).

As a general matter, monoclonal anti-3-OST antibodies may be produced by first injecting a mouse, rabbit, goat or other suitable animal with a 3-OST immunogen in a suitable carrier or diluent. As above, carrier proteins or adjuvants may be utilized and booster injections (e.g., bi- or tri-weekly over 8-10 weeks) are recommended. After allowing for development of a humoral response, the animals are sacrificed and their spleens are removed and resuspended in, for example, phosphate buffered saline (PBS). The spleen cells serve as a source of lymphocytes, some of which are producing antibody of the appropriate specificity. These cells are then fused with an immortalized cell line (e.g., myeloma), and the products of the fusion are plated into a number of tissue culture wells in the presence of a selective agent such as HAT. The wells are serially screened and replated, each time selecting cells making useful antibody. Typically, several screening and replating procedures are carried out until over 90% of the wells contain single clones which are positive for antibody production. Monoclonal antibodies produced by such clones may be purified by standard methods such as affinity chromatography using Protein A Sepharose, by ion-exchange chromatography, or by variations and combinations of these techniques.

The antibodies of the invention may be labeled or conjugated with other compounds or materials for diagnostic and/or therapeutic uses. For example, they may be coupled to radionuclides, fluorescent compounds, or enzymes for imaging or therapy, or to liposomes for the targeting of compounds contained in the liposomes to a specific tissue location.

### Assays for Drugs Which Affect 3-OST Expression

In another series of embodiments, the present invention provides assays for identifying small molecules or other compounds which are capable of inducing or inhibiting the expression of the 3-OST genes and proteins. The assays may be performed *in vitro* using non-transformed cells, established cell lines, or the transformed cells of the invention, or *in vivo* using normal non-human animals or the transgenic animal models of the invention.

In particular, the assays may detect the presence of increased or decreased expression of nucleic acids under the transcriptional control of 3-OST promoter and regulatory sequences on the basis of increased or decreased mRNA expression (using, e.g., the nucleic acid probes disclosed and enabled herein), increased or decreased

levels of protein products encoded for such nucleic acids (using, e.g., the anti-3-OST antibodies disclosed and enabled herein), or increased or decreased levels of activity of such a protein (e.g.,  $\beta$ -galactosidase or luciferase).

Thus, for example, one may culture cells known to express a particular 3-OST, or recombination modified to express at least a functional fragment or epitope of 3-OST protein under the transcriptional control of 3-OST promoter and add to the culture medium one or more test compounds. After allowing a sufficient period of time (e.g., 0-72 hours) for the compound to induce or inhibit the expression of the 3-OST, any change in levels of expression from an established baseline may be detected using any of the techniques well known in the art. Using the nucleic acid probes and/or antibodies disclosed and enabled herein, detection of changes in the expression of a 3-OST, and thus identification of the compound as an inducer or inhibitor of 3-OST expression, requires only routine experimentation. For example, one may assay for 3-OST activity by measuring the conversion of  $HS^{Inact}$  into  $HS^{Act}$  by methods known in the art (70).

In other embodiments, a recombinant assay is employed in which a reporter gene is operably joined to 3-OST promoter and regulatory sequences so as to be under the transcriptional control of these sequences. The reporter gene may be any gene which encodes a transcriptional or transitional product which is readily assayed or which has a readily determinable affect or phenotype. Preferred reporter genes are those encoding enzymes with readily detectable activity, including without limitation  $\beta$ -galactosidase, green fluorescent protein, alkaline phosphatase, or luciferase is operably joined to the 5' regulatory regions of a 3-OST gene. The 3-OST regulatory regions, may be readily isolated and cloned by one of ordinary skill in the art in light of the present disclosure of the coding regions of these genes. The reporter gene and regulatory regions are joined in-frame (or in each of the three possible reading frames) so that transcription and translation of the reporter gene may proceed under the control of the 3-OST regulatory elements. The recombinant construct may then be introduced into any appropriate host cell as described herein. The transformed cells may be grown in culture and, after establishing the baseline level of expression of the reporter gene, test compounds may be added to the medium. The ease of detection of the



expression of the reporter gene provides for a rapid, high through-put assay for the identification of inducers and inhibitors of the 3-OST gene.

Compounds identified by this method will have potential utility in modifying the expression of the 3-OST genes *in vivo*. These compounds may be further tested in the animal models disclosed and enabled herein to identify those compounds having the most potent *in vivo* effects.

#### Methods for Heparan Modification

In another aspect, the present invention provides methods for 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides in which the polysaccharides include a polysaccharide sequence of GlcA→GlcNS ±6S. These methods comprise contacting the GlcA→GlcNS ±6S-containing polysaccharide preparation with 3-OST protein in the presence of a sulfate donor under conditions which permit the 3-OST to convert the GlcA→GlcNS ±6S sequence to GlcA→GlcNS 3S ±6S. In particular embodiments, the GlcA→GlcNS ±6S sequence comprises a part of an HS<sup>act</sup> precursor sequence (i.e., GlcA→GlcNS ±6S→IdoA 2S→ GlcNS ±6S or IdoA→GlcNAc 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S) or a part of an HS<sup>inact</sup> precursor sequence (i.e., IdoA→GlcNS 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S; IdoA→GlcNAc→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S; IdoA→GlcNS→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S; IdoA→GlcNAc 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS or IdoA→GlcNS 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS). Conversion of the HS<sup>act</sup> precursor pool to HS<sup>act</sup> increases the fraction with AT-binding activity and is particularly useful in the production of anticoagulant heparan sulfate products. Thus, in another embodiment, the present invention provides for means of enriching the AT-binding fraction of a heparan sulfate pool by contacting the polysaccharide preparation with 3-OST protein in the presence of a sulfate donor under conditions which permit the 3-OST HS<sup>act</sup> conversion activity. In preferred embodiments, the sulfate donor is 3'-phospho-adenosine 5'-phosphosulfate (PAPS).

#### Methods of Partially Sequencing Complex Polysaccharides

In another aspect, the present invention provides methods for partially sequencing complex polysaccharides such as heparan sulfates (HS) or other

glycosaminoglycans (GAGs). In these methods, a pool of polysaccharides which includes sequences which may be 3-O-sulfated is contacted with a 3-OST protein in the presence of a sulfate donor (e.g., PAPS) under conditions which permit sulfation by 3-OST. The treated polysaccharides are then subjected to degradation by enzymes which degrade polysaccharides in a sequence-specific manner (e.g., polysaccharide lyases; heparinase I, II or III) and the size profile of the resulting fragments is determined. An identical pool which has not been treated with 3-OST is similarly cleaved by the same enzymes and a size profile determined. Changes in the size profiles indicate that 3-OST activity has modified the saccharide units so as to prevent (or permit) cleavage at sites which previously were (or were not) cleaved. Thus, comparison of the profiles will indicate positions at which the target sequences for 3-OST activity are present and provide a partial polysaccharide sequence.

In another embodiment, the sequence of complex polysaccharides such as HS or GAG may be partially determined using sequence specific polysaccharide affinity fractionation. To this end, 3-OST proteins which lack enzymatic function can be identified or produced (e.g., altering or deleting a portion of the catalytic ST domain by site-directed or deletion mutagenesis). These inactive forms will bind GAGs in a sequence dependent manner. For example, the 3-OST-1 protein normally, minimally, binds a GAG sequence containing GlcA-GlcNS  $\pm$ 6S. When the active site of this protein is neutralized, the  $k_d$  of the protein for these sequences will be relatively unaffected. This reagent will allow sequence-specific saccharide affinity fractionation from complex mixtures of GAGs. The purified structures can be degraded in a step-wise fashion with exolytic, endolytic enzymes and/or nitrous acid, and the resulting degradation products can be compared to standard compounds of known structure. This method will allow the quantitation and characterization of known structures contained within unknown complex polysaccharide samples.

In another embodiment, partial sequence can be obtained using the 3-OSTs of the invention or other heparan sulfate sequence specific binding ligands as protective groups prior to treating the HS or GAG with modifying agents that detectably alter the HS or GAG. Useful protective groups include catalytically inactive enzymes, chimeric enzymes and small molecule ligands with identified sequence binding specificities. The protecting group is contacted with the heparan or other

glycosaminoglycans (GAGs), and the resultant complex is treated with one or more modifying agents. Useful modifying agents include catalytically active heparan lyases, sulfotransferases, N-deacetylases, epimerases, or chimeric proteins of the invention. In embodiments where multiple protecting groups and/or modifying reagents are used in combination, the sample is first contacted with the protective group, then each modifying reagent may be with contacted with the protected polysaccharide, either simultaneously or in turn. The protective group will interfere with the ability of a chemically modifying agent to interact with, attach to and/or cleave specific GAG sequence motifs. The sample can then be analyzed for ligand-specific protection and/or cleavage to elucidate the sequence of the original GAG using separation and/or quantitation using methods known in the art.

In some embodiments, as a preliminary step, full length heparans and GAG oligomers can be fractionated over an immobilized affinity ligand immobilized at their reducing ends *via* hydrazide chemistry. The fraction of GAG captured by the immobile phase permits a quantitation of the mass or total percent of the target sequence (out of total GAG.) Thus, unique heparan or other GAG structures may be concentrated and/or specifically eluted for further analysis.

One useful method for the detection binding is the Biomolecular Interaction Assay or "BIAcore" system developed by Pharmacia Biosensor and described in the manufacturer's protocol (LKB Pharmacia, Sweden). In light of the present disclosure, one of ordinary skill in the art is now enabled to employ this system, or a substantial equivalent, to identify proteins or other compounds having sequence-specific HS or GAG binding capacity, or HS or GAGs sequences having 3-OST binding capacity. Such systems utilize surface plasmon resonance, an optical phenomenon that detects changes in refractive indices. A sample of interest is passed over an immobilized ligand (e.g., a 3-OST fusion protein or specific GAG) and binding interactions are registered as changes in the refractive index.

### Examples

#### Cell Lines and Cell Culture

The clonal L cell line LTA (35, 41), the generation of clone 33, an LTA transfectant that over-expresses the ryudocan<sub>12CA5</sub> cDNA (33), a rapidly growing

revertant of clone 33, L-33<sup>+</sup> (26), and RFPEC, an immortalized line derived from rat fat-pad endothelial cells (8) have previously been described. Primary mouse neonatal endothelial cells from the cardiac microvasculature of day 3-5 neonates (CME cells) (from Dr. Jay Edelberg, MIT/Beth Israel Hospital) and COS-7 cells (ATCC) were employed. Primary human umbilical cells (HUVEC) were maintained according to the supplier's (Clonetics Inc.) protocol. Unless otherwise stated, all cell lines were maintained in logarithmic growth by subculturing biweekly in Dulbecco's modified Eagle medium (Life Technologies, Inc.) containing 10% fetal bovine serum, 100 µg/ml streptomycin, and 100 units/ml penicillin at 37 °C under 5% CO<sub>2</sub> humidified atmosphere, as previously described (42). Exponentially growing cultures were generated by inoculating 54,000 cells/cm<sup>2</sup> and incubating for two days, whereas post-confluent cultures were produced by inoculating 250,000 cells/cm<sup>2</sup> and allowing growth for 10 days with medium exchanges on days 4, 7, 8, and 9.

#### Peptide Purification and Sequencing

The purification of mouse 3-OST-1 from L-33<sup>+</sup> has been previously described (26) and the final step 4 product was concentrated by reverse phase chromatography on a HP 1090 M system (Hewlett Packard) equipped with a C4 reverse phase HPLC column (250 x 2.1 mm, 300 Å pore size, 5 µm particle size) (Vydac, number 214TP52) equilibrated in 1.6% acetonitrile (v/v), 0.1% TFA (v/v). After application of sample, the reverse phase matrix was washed with 60% acetonitrile, 0.1% TFA, and bound species were eluted with 78.4% acetonitrile, 0.1% TFA. Samples of 1.5 or 3 µg, from two independent purifications, were digested with 0.15 or 0.3 µg, respectively, of endopeptidase Lys-C (Waco) in a reaction volume of 100 µl containing 1% RTX100 (Calbiochem), 10% acetonitrile and 100 mM Tris-HCl pH 8.0, at 37 °C for ~16 h (43). Digestion products were chromatographed on an HP 1090 M system (Hewlett Packard) equipped with the above described C4 reverse phase HPLC column equilibrated in 98% Buffer A (0.1% TFA (v/v))/2% Buffer B (80% acetonitrile (v/v)/0.85% TFA (v/v)). After application of digestion products, the reverse phase matrix was washed with 98% Buffer A/2% Buffer B, and bound species were eluted with linear gradients of Buffer B increasing to 37.5% over 60 min, to 75% over 30 min, and to 98% over 15 min (44). The eluate was monitored for absorbance at 210 and 280 nm, peptide peaks were individually collected and analyzed with a

model 477A/120A Protein Sequenator (Applied Biosystems). In addition, the NH<sub>2</sub>-terminal sequence of 1 µg of concentrated 3-OST-1 sample was directly determined.

#### Isolation of Mouse 3-OST-1 Clones

Isolation of Cytoplasmic and Poly(A)<sup>+</sup> RNA. Cytoplasmic RNA (17.5 mg) was isolated from post-confluent cultures of LTA cells (12 flasks of 175 cm<sup>2</sup>, ~1.6 x 10<sup>9</sup> cells) by a modification of the procedure of Favalaro (45). Monolayers were twice washed with PBS, cells were recovered by trypsinization and centrifugation (1000 x g for 2 min), and cell pellets were washed by resuspension in PBS followed by centrifugation (1300 x g for 4 min). Cells were lysed by vortexing for 30 sec in 12 ml of ice cold 50 mM Tris, pH 7.4, 140 mM NaCl, 5 mM EDTA, 1 % Triton X-100, 5 mM vanadium ribonucleoside complexes (Life Sciences Technologies), samples were incubated on ice for 10 min and then vortexed for 1 min. Nuclei were pelleted by centrifugation at 6000 x g for 10 min, the supernatant was mixed with an equal volume of 200 mM Tris, pH 7.4, 300 mM NaCl, 2% SDS, 25 mM EDTA, containing 200 µg/ml of proteinase K (Boehringer Mannheim), and the mixture was incubated at 65 °C for 2 hr. Samples were extracted twice against an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1), the aqueous phase was combined with 0.7 volumes of isopropanol, cytoplasmic RNA was pelleted by centrifugation at 3500 x g for 10 min, and was resuspended in 3.6 ml of 10 mM Tris, pH 7.4, 1 mM EDTA. Poly(A)<sup>+</sup> RNA (59 µg) was isolated from 16 mg of cytoplasmic RNA by two sequential purifications against 100 mg of oligo(dT) cellulose (Life Sciences Technologies, #15939-010) according to the manufacturer's specifications except that binding and wash buffers contained 0.1 % SDS and LiCl was substituted for NaCl. The final eluate (1.5 ml) was extracted against 1.5 ml of phenol/chloroform/isoamyl alcohol (25:24:1), the aqueous phase was then adjusted to 100 mM LiCl and 260 mM NaCl, an equal volume of isopropanol was added, the mixture was centrifuged at 15,000 x g for 30 min and the poly(A)<sup>+</sup> RNA pellet was recovered in 40 µl of diethyl pyrocarbonate treated water.

PCR Cloning and Generation of a Mouse 3-OST-1 Probe. Degenerate PCR primers 1S, 2S, 2A, and 3A (described in Shworak et al. (1997) J. Biol. Chem. 272, in press) were obtained from Bio Synthesis. First strand cDNA was generated in a 50 µl volume from 5 µg of LTA poly(A)<sup>+</sup> RNA primed with oligo(dT) using an RT-PCR kit

(Stratagene, La Jolla, CA) according to the manufacturer's specifications. Touchdown PCR (46, 47) reactions (50  $\mu$ l) contained 1  $\mu$ l of first strand cDNA, 25 pmol of each primer, 0.25  $\mu$ l of AmpliTaq Gold (Perkin Elmer), 200  $\mu$ M of each dNTP and 1 x GeneAmp PCR buffer. Two distinct sets of touchdown PCR conditions were required to obtain optimal yields of product. For amplification with primers 1S and 2A, reactions were heated to 95  $^{\circ}$ C for 9 min, subjected to 20 cycles of 94  $^{\circ}$ C for 30 sec, and 68  $^{\circ}$ C for 1 min with a 0.5  $^{\circ}$ C reduction per cycle, followed by 20 cycles of 94  $^{\circ}$ C for 30 sec, 58  $^{\circ}$ C for 30 sec with a 0.5  $^{\circ}$ C reduction per cycle, and 75  $^{\circ}$ C for 30 sec, then 15 cycles of 94  $^{\circ}$ C for 30 sec, 55  $^{\circ}$ C for 10 sec, and ramping to 75  $^{\circ}$ C over 50 sec. Alternatively, for amplification with primers 1S and 3A or primers 2S and 3A, reactions were heated to 95  $^{\circ}$ C for 4 min, subjected to 47 cycles of 95  $^{\circ}$ C for 30 sec, and 69.5  $^{\circ}$ C for 2 min with 0.2  $^{\circ}$ C and 1 sec reductions per cycle, followed by 25 cycles of 95  $^{\circ}$ C for 30 sec, 60  $^{\circ}$ C for 15 sec, and ramping to 75  $^{\circ}$ C over 1 min. Amplification products were purified as the retentate from centrifugal ultrafiltration against a 30,000 molecular weight cutoff membrane (Millipore, # SK1P343JO), then 200 ng of DNA was end polished with *Pfu* DNA polymerase and subcloned into pCR-Script Amp SK(+) (Stratagene, La Jolla, CA, #211188) according to the manufacturer's specifications. A resulting plasmid, pNWS182, contained the 1S/3A amplification product of 779 bp which was released by digestion with *Eco*RI and *Sac*II, and isolated by low melting point agarose gel electrophoresis. A  $^{32}$ P-labeled primer extension probe was then generated with a random primer labeling kit (Stratagene, La Jolla, CA, # 300385) by replacing the random primers with 5  $\mu$ M of primer 3A.

#### Construction and Screening of an L Cell cDNA Library.

Using the manufacturer's recommended conditions, an oligo(dT)-primed  $\lambda$  Zap Express cDNA library (Stratagene, La Jolla, CA, # 200451) was generated from 5  $\mu$ g of LTA poly(A)<sup>+</sup> RNA which had been pretreated with methylmercury hydroxide. About  $1.5 \times 10^6$  primary recombinants were plaque amplified by infection into *E. coli* XL1-Blue MRF'. From the amplified library,  $1.3 \times 10^6$  plaques were transferred to Colony/Plaque Screen (Du Pont-New England Nuclear) and screened with the above described  $^{32}$ P-labeled probe specific for 3-OST-1. Hybridizations were performed at

42 °C in 1.7 x SSC, 8.3% dextran sulfate, 42% formamide, 0.8% SDS and filters were washed twice with 2 x SSC, 1% SDS for 30 min at 65 °C. Positive clones were plaque purified and then *in vivo* excised into pBK-CMV based phagemids by infection with ExAssist helper phage followed by transduction of filamentous phage particles into *E. coli* XL0LR.

#### Isolation of Human 3-OST-1 cDNA Clones

The National Center for Biotechnology Information data bank of I.M.A.G.E. Consortium (LLNL) expressed sequence tag cDNA clones (48) was probed with the deduced mouse 3-OST-1 amino acid sequence to reveal three partial length species.

I.M.A.G.E. Consortium CloneID 220372 (accession numbers H86812 and H86876) was from the retinal library of Soares (N2b4HR), whereas clones 301725 (accession numbers N90867 and W16558) and 301726 (accession numbers N90856 and W16555) were from the fetal lung library of Soares (NbHL19W ) and were obtained from the TIGR/ATCC Special Collection (ATCC). The *EcoRI/NotI* insert of clone 220372 was <sup>32</sup>P labeled by random priming and used to screen 5 x 10<sup>5</sup> plaques from a

λ TriplEx Brain cDNA library (Clontech, Palo Alto, CA), as described above. Positive plaques were purified, TriplEx based plasmids were *in vivo* excised according to the manufacturer's protocol, and were sequenced as described below.

#### Characterization of Mouse and Human 3-OST-1 cDNA Clones

The 5' and 3' regions of all partial and full length clones were enzymatically sequenced from flanking primer sites of the respective cloning vectors. For full length clones the remaining sequence of both strands was obtained with internally priming oligonucleotides. Automated fluorescence sequencing was performed with Perkin Elmer Applied Biosystems Models 373A and 477 DNA sequencers. Each reaction typically yielded 400 to 600 bases of high quality sequence. cDNA sequence files were aligned and compiled with the program Sequencher 3.0 (Gene Codes Corp.). All additional manipulations were performed with the University of Wisconsin Genetics Computer Group sequence analysis software package. Sequence comparison searches were performed on the databases of GenBank, EMBL, DDBJ, PDB, SwissProt, PIR, and dbEST.

#### Expression of 3-OST-1 cDNAs

Construction of Expression Plasmids. The plasmid pCMV-3-OST contains the mouse 3-OST-1 cDNA, an *EcoRI/XhoI* fragment from pNWS228, inserted between the CMV promoter and the bovine growth hormone polyadenylation signal of *EcoRI/XhoI* digested and phosphatase treated pcDNA3 (Invitrogen). The plasmid pCMV-ProA3-OST is of similar structure, except the first 26 amino acid of 3-OST-1 are replaced with 291 amino acids encoding a fusion protein of the transin leader sequence followed by Protein A and a factor Xa cleavage site. pCMV-ProA-3-OST was generated by ligating a *BamHI/SmaI* fragment containing the Protein A region from pRK5F10PROTA (49), and an *XmaI* (end-filled with T<sub>4</sub> polymerase)/*XhoI* fragment containing most of the mouse 3-OST-1 cDNA from pNWS228, into *BamHI/XhoI* digested and phosphatase treated pcDNA3 (Invitrogen). The *in vitro* transcription plasmid, pNWS237, contains a T3 promoter site 5' of the human 3-OST-1 cDNA and was constructed by inserting complementary oligonucleotides (Bio Synthesis) into the *EcoRI* site of the TriplEx based plasmid, pJL30.

Transient Expression of the Mouse 3-OST-1 cDNA in COS-7 Cells. For each expression construct, three 175 cm<sup>2</sup> flasks were seeded with 3.6 x 10<sup>6</sup> COS-7 cells, 6 h later the medium was exchanged with DMEM containing 10% Nu-Serum (Life Technologies, Inc.) with 100 µg/ml streptomycin and 100 units/ml penicillin, and cells were grown for an additional day. Monolayers were washed with PBS then incubated at 37 °C for 2.5 h with 10 ml/flask of freshly prepared DMEM containing 235 µg/ml DEAE-dextran (M.W. 500,000, Pharmacia), 9.5 mM Tris-HCl, pH 7.4, 0.9 mM chloroquine-diphosphate (Sigma), and 3 µg/ml of the appropriate pcDNA3 based expression plasmid. Monolayers were then exposed to freshly prepared 10% DMSO in PBS for 1.5 min, washed twice with nonsupplemented DMEM, fed 30 ml/flask of DMEM containing 10% fetal bovine serum, 100 µg/ml streptomycin, and 100 units/ml penicillin, and cells were grown for an additional day. Monolayers were washed with PBS, then cells were grown in 40 ml/flask Serum-Free Medium (DMEM containing 25 mM HEPES, pH 8.0, 1% Nutridoma SP (Boehringer Mannheim) (v/v), an additional 2 mM glutamine, 10 ng/ml biotin (Pierce), 100 µg/ml streptomycin, 100 units/ml penicillin, and 1 x of a previously described Trace Metal Mix (26)) for 24 h. COS-cell conditioned Serum-Free Medium was harvested, debris was removed by centrifugation at 1,000 x g for 10 min followed by filtration through a 0.45 µm



membrane, then samples were either immediately processed or were snap frozen with liquid nitrogen and stored at -80 °C. Occasionally, conditioned medium from a second incubation of 8-24 h was also collected.

#### Purification of Wild-type and Protein A Tagged Mouse Recombinant 3-OST-

1. Wild-type mouse recombinantly expressed 3-OST-1 enzyme (r3-OST-1) was purified, at 4 °C, from 240 ml of freshly generated Serum-Free Medium conditioned by COS-7 cells transfected with pCMV-3-OST. The medium was adjusted to pH 8.0, mixed with an equal volume 2% glycerol, then loaded (25 ml/h) onto a heparin-AF Toyopearl-650M column (0.8 x 5.7 cm) (TosoHaas, Montgomeryville, PA) equilibrated in 50 mM NaCl, 10 mM Tris-HCl, pH 8.0, 1% glycerol (v/v) (Buffer C). The column was washed with 20 ml of Buffer C at a flow rate of 0.8 ml/min, then with 20 ml of 150 mM NaCl, 10 mM Tris-HCl, pH 8.0, 1% glycerol (v/v) at a flow rate of 0.5 ml/min, and protein was eluted at a flow rate of 0.25 ml/min with a 20 ml linear NaCl gradient extending from 150 mM to 750 mM NaCl in Buffer C. The fractions exhibiting HS<sup>act</sup> conversion activity (approximately 4 ml) were pooled, brought to a final concentration of 0.6% CHAPS (w/v) (Sigma) and dialyzed for 16 h against 4 l of 25 mM MOPS (3-[N-morpholino] propanesulfonic acid) (Sigma), pH 7.0, 1% glycerol (v/v), 0.6% CHAPS (w/v) (MCG buffer ) containing 50 mM NaCl. The dialysate was applied to a 3',5'-ADP-agarose column (0.8 x 1.2 cm, 3.7 mmol of 3',5'-ADP/ml of gel) (Sigma ) and eluted as previously described (26). The fractions containing HS<sup>act</sup> conversion activity were pooled (approximately 4 ml), aliquoted, frozen in liquid nitrogen and stored at -80 °C.

Protein A tagged mouse r3-OST-1 was purified, at 4 °C, from 155 ml of previously frozen Serum-Free Medium conditioned by COS-7 cells transfected with pCMV-ProA3-OST. IgG agarose beads (310 µl of a 50/50 slurry; Sigma) were gently stirred with the conditioned medium for 3h, recovered by centrifugation at 2,000 x g for 10 min, and washed twice with 1 ml of MCG containing 250 mM NaCl to remove nonspecifically bound protein. Protein A fusion-protein was eluted from the beads with two sequential 30 min incubations in 100 µl of 50 mM sodium acetate, pH 4.5, 150 mM NaCl, 0.6% CHAPS and 1% glycerol. The pooled eluates were combined with an equal volume of 500 mM MOPS, pH 7.0, 0.6% CHAPS, and 1% glycerol, then aliquoted, frozen in liquid nitrogen and stored at -80 °C.

## Retroviral Transduction of CHO and MNE Cells with 3-OST-1

Plasmid retrovirus vector construction. A retroviral transduction system was used to transduce CHO cells and mouse neonatal endothelial (MNE) cells. This system may serve as a model for in vivo transduction for use in gene therapy.

5 The retrovirus backbone plasmid pMSCV-PGK-EGFP is a derivative of pMSCVpac a (Dr. Robert Hawley University of Toronto.) The puromycin acetyl transferase gene cassette in pMSCVpac was removed and replaced with an Enhanced GFP (Dr. David Baltimore MIT). The pMSCV-PGK-GFP vector was assembled by digestion of the plasmid with HindIII and ClaI, followed by treatment with Klenow  
10 fragment. The EGFP cistron 720 bp fragment was derived from the digestion of pMSCV-EGFPpac with EcoRI, and blunting with the Klenow fragment. The EGFP blunt-ended fragment was then ligated into the blunt-ended pMSCV vector. The resulting plasmids were tested for proper orientation by restriction analysis. The reporter virus, pMSCVPLAP, is designed to express the wild type human placental  
15 alkaline phosphatase (PLAP) transcribed from the 5' LTR. pMSCV-SEAP-PGK-EGFP was made by cloning the secreted alkaline phosphatase (SEAP) BglIII and HpaI 1.723 kb fragment from pSEAP2-basic plasmid (Clontech, Palo Alto, CA) into the BglIII and HpaI cut pMSCV-PGKEGFP vector. pCMV3-OST was digested with BglIII and XhoI to release the wild type mouse 3-OST-1 cDNA. The 1.623 kb 3-OST-  
20 1 cDNA fragment was cloned into the BglIII and XhoI sites in pMSCV-PGK-EGFP. The occurrence of the insert of interest present in the correct orientation was ascertained by restriction analysis. All plasmid DNA prepared for transfection was made with the Invitrogen SNAP-MIDI kits according to the manufacturer's directions.

Cells and cell culture. Dulbecco's modified Eagle medium (DMEM), F-12  
25 Ham's medium and penicillin/streptomycin, 0.25% trypsin, 10 mM EDTA, were obtained from Life Technologies, Inc., GIBCO-BRL (Gaithersburg, MD). The PHOENIX ecotropic retroviral packaging cell line (ATCC #SD 3444) was grown in DMEM, 10% heat-treated fetal bovine serum (FBS) (JRH Biosciences, Lenexa, KS), 100 units/ml penicillin, 100 µg/ml. PHOENIX cells were subcultured three times  
30 weekly at a split ratio of approximately 1:8 in a 37 °C humidified, 5.0% CO<sub>2</sub> incubator. CHOK1 ATCC CCL 61 cells (CHO) were grown in F-12 medium supplemented with 10% fetal bovine serum, and 100 units/ml penicillin, 100 µg/ml in

a 37 °C humidified, 5.0% CO<sub>2</sub> incubator. CHO cells were subcultured three times weekly at a split ratio of approximately 1:4 in a 37 °C humidified, 5.0% CO<sub>2</sub> incubator. 1 x 10<sup>6</sup> CHO cells were transfected with 10 µg of pcB7-ECOTROPIC (generous gift of Dr. Harvey Lodish) by the standard calcium phosphate precipitation technique. Plasmid pcB7-ECOTROPIC expresses the MCAT1 gene (ecotropic retrovirus receptor cDNA) and hygromycin resistance gene transcribed from separate constitutive promoters. The transfected cells were selected for hygromycin resistance in 200 µg/ml hygromycin (Life Technologies). The stable, hygromycin-resistant clones were assayed for their ability to take up and express reporter virus (MSCVPLAP). Fixation and staining for cell-bound alkaline phosphatase was performed by standard techniques. CHO clone 4B was chosen because it transduced most efficiently at the highest dilution tested (i.e., 1:10,000), and was expanded for further analysis. Transduction of CHO4B with ecotropic retroviruses is equal to that achievable with NIH3T3 cells. Low passage number (passage 2-5), primary mouse neonatal cardiac endothelial cells (MNE) were prepared by standard techniques. MNE cells were cultured in a 1:1 vol./vol. admixture of EGM:EGM-2 (CLONETICS) in a 37 °C, humidified, 5.0% CO<sub>2</sub> incubator. MNE cells were subcultured once weekly at a split ratio of approximately 1:3 in a 37 °C, humidified, 5.0% CO<sub>2</sub> incubator.

Northern blot analysis. Total RNA was prepared from confluent T-80 flasks of each of the transduced and untransduced cells using the QIAGEN RNAeasy kit with QIASHREDDER. 10 µg of total cellular RNA was denatured and resolved by electrophoresis in a 1.5% agarose gel, and then blotted onto GENE-Screen+ (DuPont NEN) with 2X SSPE. The membrane was then UV cross-linked using a STRATALinker. <sup>32</sup>P-radiolabeled cDNA probes were prepared from the fragments of DNA used for cloning the mouse 3-OST-1 and SEAP as described above. Radiolabeled probes were prepared using 25ng of each template and the Amersham Megaprime kit, and α <sup>32</sup>P dCTP from DuPont NEN according to the manufacturer's directions. Hybridizations were performed in sealable plastic bags at 68 °C with 1 x 10<sup>6</sup> cpm of probe/ml in 10 ml of QUICKHYB (Stratagene, La Jolla, CA), following the manufacturer's instructions. Post-hybridization washes were: once for 15 minutes

in 1X SSPE, 1.0% SDS at 45 °C; and then twice for 15 minutes each in 0.2X SSPE, 0.5% SDS 650C. After washing, the blots were briefly air dried, placed in sealable plastic bags then exposed to Kodak XAR-MS film with intensifying screens at -80 °C for from overnight to five days. Quantitation of hybridizing signal intensity was performed using a Betascope 603 blot analyzer. Transcripts derived from the 5' LTR of these engineered proviruses are large (ca. 7 kb). Since they are large, have multiple sites of transcriptional initiation provirus (5' LTR and pgk promoters), and the 3-OST-1 construct has more than one poly(A) addition signal, bona-fide hybridizable mRNA will appear as different sizes in northern blot analysis. The total amount of hybridizing material detected, per sample lane, with any one probe was used to calculate and compare mRNA expression levels.

Virion production. Virions were produced by programming ecotropic PHOENIX packaging cells with recombinant provirus plasmids using the calcium phosphate transfection technique. 10 µg/well of each recombinant retroviral construct plasmid was transfected via calcium precipitation with an overnight incubation period. Following the precipitation step, the cells were re-fed with 2 ml/well of fresh DMEM and incubated overnight. Each 2 ml of viral supernatant was collected and flash-frozen in liquid nitrogen and stored at -80 °C, or used directly after a low-speed centrifugation.

Transduction protocol. Target cells were trypsinized, counted with a Coulter cell counter and then plated at 150,000 cells (NIH 3T3/CHO4B) or 50,000 cells (MNE) per well of a cluster-6 well plate. 24 hours later, target cells (<70% confluent) were incubated overnight with viral supernatants containing as adjuvants either 5 µg/ml polybrene for NIH3T3/CHO4B or 25 µg/ml DEAE-dextran (Pharmacia) for MNE. After 12 hours of virus exposure, the growth media was replaced. CHO cells destined for FACS sorting were exposed to recombinant retrovirus two times at a multiplicity of infection (MOI) of 0.3. MNE cells were transduced one time for 12 hours at an MOI of 0.74 for recombinant 3-OST-1 virus and 0.72 for recombinant SEAP virus. Transduced cells were allowed to incubate in fresh growth medium for 48 hours prior to FACS to allow for maximum proviral expression. Recombinant virus titers ranged from  $1 \times 10^5$  -  $2 \times 10^6$  infectious particles per ml as measured with either NIH3T3 or CHO4B cells using FACS analysis scoring for EGFP positive cells.

Virus titers were reduced approximately eight to ten-fold on primary MNE cells relative to NIH3T3.

#### Cell-Free Synthesis of Mouse and Human r3-OST-1.

Synthetic capped mouse and human 3-OST-1 mRNAs were generated from  
5 *NotI* linearized pNWS228 and *HinDIII* linearized pNWS237, respectively, using T<sub>3</sub>  
polymerase and m<sup>7</sup>G(5')ppp(5')G, as previously described (50). Unlabeled *in vitro*  
translation reactions (25 µl) contained 0.25 µg of synthetic mRNA, 1.8 µl canine  
pancreatic microsomal membranes (Promega), 0.5 µl each of Amino Acid Mixture  
Minus Leucine and Amino Acid Mixture Minus Methionine, and were performed with  
10 nuclease-treated reticulocyte lysate (Promega), according to the manufacturer's  
specifications.

Measurement of HS<sup>act</sup> Conversion Activity. The HS<sup>act</sup> conversion activity, a  
3-OST-1 catalyzed reaction which requires unlabeled PAPS to convert <sup>35</sup>S-HS<sup>inact</sup> into  
<sup>35</sup>S-HS<sup>act</sup>, of crude and purified r3-OST-1 samples was determined by comparison  
15 against a standard curve generated with 1 to 32 units of previously purified native 3-  
OST-1, as previously described (26). The <sup>35</sup>S-HS<sup>inact</sup> substrate was purified from  
metabolically labeled cell surface HS of exponentially growing clone 33 cells, as  
previously described (35).

#### Identification of Enzymatic Reaction Products

20 <sup>35</sup>S-labeling of HS by r3-OST-1. <sup>35</sup>S-labeled HS was generated by incubating  
the various forms of r3-OST-1 with [<sup>35</sup>S]PAPS and unlabeled HS<sup>inact</sup>, which were  
prepared as previously described (26, 35). Wild-type and Protein A tagged r3-OST-1  
(2500 units of HS<sup>act</sup> conversion activity) purified from COS cell conditioned medium,  
were incubated in a 500 µl reaction mixture, as previously described (26), for 2 h at  
25 37 °C and <sup>35</sup>S-labeled polysaccharides were purified by DEAE-Sepharose  
chromatography as previously described (26). For cell-free synthesized r3-OST-1,  
<sup>35</sup>S-labeling of HS was performed in a reticulocyte lysate based reaction mixture (35)  
except that 100 µl reactions contained 100 to 300 units of *in vitro* translated r3-OST,  
180 nM unlabeled HS<sup>inact</sup>, 5 µM PAPS (60 x10<sup>6</sup> cpm) and samples were incubated at  
30 37 °C for 2 h. The reaction was quenched by the addition of 300 µl of 267 mM NaCl,  
13.3 µg/ml glycogen and extraction against 600 µl of phenol/chloroform/isoamyl

alcohol (25:24:1). <sup>35</sup>S-labeled GAGs were ethanol precipitated (35) and then isolated by DEAE chromatography as previously described (26).

Identification of the Site of Sulfation on HS<sup>act</sup> and HS<sup>inact</sup>. The DEAE eluates containing <sup>35</sup>S-labeled polysaccharide were vacuum concentrated to 1/5 volume, then  
5 desalted at a flow rate of 0.9 ml/min on TSK G3000 PW<sub>XL</sub> (0.78 x 30 cm) and TSK G2500 PW<sub>XL</sub> (0.78 x 30 cm) (TosoHaas) columns connected in series equilibrated in 0.1 M ammonium bicarbonate. The desalted product was then affinity fractionated using AT/ConA gel to obtain HS<sup>act</sup> and HS<sup>inact</sup> as described previously (26). Analysis of labeled products by treatment with GAG lyases and low pH nitrous acid were  
10 performed as previously described (42). In addition, the HS<sup>act</sup> and HS<sup>inact</sup> samples were each subjected to hydrazinolysis, high pH nitrous acid (pH 5.5), low pH nitrous acid (pH 1.5), and sodium borohydride reduction with the resultant disaccharides characterized on reverse phase ion pairing HPLC (RPIP-HPLC) as previously reported (33, 34). The identification of [<sup>35</sup>S]GlcA→AMN-3-O-SO<sub>3</sub> and [<sup>35</sup>S]GlcA→AMN-  
15 3,6-O-(SO<sub>3</sub>)<sub>2</sub> was confirmed by co-chromatography on RPIC-HPLC with the appropriate <sup>3</sup>H-labeled disaccharide standards, as described in prior publications (33,34).

#### Northern Blot Analysis

Total RNA from RFPEC and primary mouse CME cells was isolated by the  
20 method of Chomczynski and Sacchi (51), whereas poly(A)<sup>+</sup> RNA was isolated from HUVEC cells as described above for LTA cells. Total RNA from the mast cell line CLMC/C57.1 (C57.1) (52) was from Dr. Stephen J. Galli (Beth Israel Hospital). Samples were resolved on 1.2% formaldehyde-agarose gels and subjected to Northern blot analysis as previously described (50). Mouse and human samples were  
25 hybridized with mouse or human probes, respectively, and washed as described for library screening, above, except hybridizations were performed at 60 °C.

#### Peptide Sequencing and PCR Generation of a Mouse 3-O-Sulfotransferase-1 (3-OST-

##### 1) Probe

The information necessary for the molecular cloning of mouse heparan sulfate  
30 D-glucosaminyl 3-O-sulfotransferase-1 (3-OST-1) was obtained by sequencing the amino terminus and Lys-C generated peptides of the enzyme that we had previously purified from large quantities of serum-free tissue culture medium conditioned by an

L cell line (26). These studies established the structures of 14 partially overlapping peptides which encompass 185 amino acid residues. Degenerate PCR primers were synthesized based on the sequence of the amino terminus (primer 1S) and two endopeptidase derived fragments (primers 2S, 2A, and 3A). When PCR was performed on an LTA first strand cDNA template, products of about 210 (primers 1S/2A) and 780 (primers 1S/3A) and 610 (primers 2S/3A) bp were obtained, which suggests that all of the primer sites are contained within a single cDNA. To confirm this supposition, the two largest fragments were cloned into pCR-Script Amp SK(+) and inserts were sequenced, which revealed that the 1S/3A product is 779 bp and contains the 611 bp 2S/3A product. The 779 bp insert encodes 12 of the sequenced peptide fragments and so was <sup>32</sup>P-labeled, as described above, and used as a probe for cDNA library screening.

#### Isolation and Characterization of Mouse 3-OST-1 cDNAs

An amplified  $\lambda$  Zap Express LTA cDNA library of  $1.5 \times 10^6$  primary recombinants was constructed and  $1.3 \times 10^6$  plaques were screened with the above described probe, which revealed 40 positives that were plaque purified and *in vivo* excised into plasmids. The cDNA inserts of each plasmid were characterized to eliminate duplicated recombinants due to library amplification. Size was determined by liberating cDNA inserts with digestion at flanking *EcoRI* and *XhoI* restriction sites followed by agarose gel electrophoresis; furthermore, the sequence at both ends of each insert was obtained from flanking vector primer sites. This analysis revealed 25 unique primary recombinants which predominantly contained inserts of approximately 1.7, 2.3, or 3.3 kb. These different species were considered to reflect natural size variants of the mouse message since northern blots of LTA poly(A)<sup>+</sup> RNA hybridized with 3-OST-1 probe revealed the same three size categories of message. The complete sequencing of 9 distinct primary recombinants, at least 2 from each size category, in conjunction with the partial sequencing of the remaining 16 clones showed that the size variants result from differences in the length of 5' untranslated region due to the insertion of 0-1629 bp at a single common internal point, the splice variant site. Most importantly, all clones shared identical protein coding regions and, therefore, the characterization and analysis of only the shortest species, the Class 1 cDNA, which lacks additional sequence at the splice variant site, is described below.

Sequence data was obtained from 2 essentially full length Class 1 cDNAs, and 5 partial length cDNAs to create a composite cDNA structure of 1685 bp (SEQ ID NO: 1), excluding the 3' poly(A) tract. The 5' untranslated region is 322 bp with the splice variant site occurring between nucleotides 216 and 217. This region contains 6 ATG sites which do not conform to consensus initiation sites (53) and are followed by near in-frame termination codons. An open reading frame of 933 bp begins at position 323 with the first consensus initiation ATG (a purine occurs at -3) (53). The length of the 3' untranslated region from all of the cDNA clones analyzed ranged from 301-430 bp. Within this terminal 129 bp, 5 distinct polyadenylation sites were observed and 13-18 bp upstream from each site is a variant of the consensus polyadenylation signal. Poly(A) tails were most frequently observed at the first site (position 1556, ~50% of clones).

#### Isolation and Characterization of Human 3-OST-1 cDNAs

Three clones containing partial length human 3-OST-1 cDNAs were identified by EST database searching (48) and were obtained from the TIGR/ATCC Special Collection, as described above. Sequencing of the insert ends revealed the clones to be essentially equivalent, as each contained the same 947 bp region of the human 3-OST-1 cDNA. The insert of I.M.A.G.E. Consortium CloneID 220372 was <sup>32</sup>P-labeled and used to screen 5 x 10<sup>5</sup> plaques from a λ TriplEx Brain cDNA library. Three positives were identified and isolated as TriplEx plasmids, and the largest cDNA 1.3 kb was sequenced completely.

The nucleic acid sequence of mouse and human 3-OST-1 cDNAs are ~85% identical. The largest isolated human clone contains 118 bp of 5' untranslated region with 2 nonconsensus ATG sites. The sequences of human and mouse cDNAs flanking the splice variant site on the 5' limit are distinct (positions 211-216 of SEQ ID NO: 1 and positions 5-10 of SEQ ID NO: 3), but on the 3' limit are identical (positions 217-222 of SEQ ID NO: 1 and positions 11-16 of SEQ ID NO: 3), which raises the possibility that human 3-OST-1 mRNA may also exhibit 5' splice variants. The first consensus ATG (with a purine occurring at -3 and a G at +4) (53) initiates an open reading frame of 921 bp. For all 4 human cDNA clones examined, only a single polyadenylation site was observed resulting in a 3' untranslated region of 266 bp, which is 26 bp less than the most frequently observed 3' limit for the mouse cDNAs.



### Predicted Protein Structures of Mouse and Human 3-OST-1

The mouse and human cDNAs encode novel 311 and 307 amino acid proteins of 35,876 and 35,750 daltons, respectively, that exhibit 93% similarity. The deduced mouse primary structure contains regions corresponding to all 13 sequenced peptides and the amino terminus. For both types of 3-OST-1, the encoded protein is predicted to be an intraluminal resident. Kyte-Doolittle hydropathy analysis reveals only a single major hydrophobic region which begins at the amino terminus and lacks sufficient length for a membrane spanning domain. Moreover, the hydrophobic region differs from a membrane anchor in that it contains two glutamine residues and is not flanked by cationic residues. Thus, the above stretch of 18 residues constitutes a hydrophobic leader signal, and this region is followed by a signal peptidase cleavage site between amino acids 20 and 21, as determined by the method of von Heijne (54). The possibility of signal peptidase cleavage is supported by the amino-terminal analysis of mouse 3-OST-1, which began with His<sup>21</sup>. Given that heparan biosynthesis is considered to occur in the *trans*-Golgi, the above data suggest that the 3-OST-1 is an intraluminal enzyme. Just past the signal peptidase cleavage site, the mouse 3-OST-1 contains an extra 4 residues (Asp<sup>24</sup>-Pro<sup>25</sup>-Gly<sup>26</sup>-Pro<sup>27</sup>) not found in the human form. Both 3-OST-1 proteins exhibit 5 potential *N*-glycosylation sites which account for the apparent discrepancy between the molecular weights of the predicted amino terminus trimmed enzyme (~34 kDa) and the previously purified enzyme (a broad band of 46 kDa was observed on SDS-PAGE) (26). Only two cysteine residues are present, and these closely spaced residues are likely to form a disulfide bond which generates a peptide loop of 10 amino acids. Interestingly, the carboxy 140 residue region is extremely basic (25% H, K, R; 12% E, D); however, this region does not exhibit previously recognized heparin binding motifs.

### Recombinant Expression of Mouse and Human 3-OST-1 Enzyme (r3-OST-1)

Three distinct expression approaches were employed to confirm that the isolated cDNAs encode 3-OST-1 enzyme. The resulting recombinantly expressed 3-OST-1 enzyme was designated as r3-OST-1, to distinguish this form from the previously purified native 3-OST-1 enzyme. First, the vector pCMV-3-OST (a pcDNA3 derivative in which the CMV promoter transcribes the mouse 3-OST-1 cDNA) was transiently expressed in COS-7 cells and the resulting level of HS<sup>act</sup>

conversion activity accumulated in Serum-Free Medium over 32 h was measured, as described above.  $\text{HS}^{\text{act}}$  conversion activity is a 3-OST-1 catalyzed reaction which requires unlabeled PAPS to convert  $^{35}\text{S-HS}^{\text{inact}}$  into  $^{35}\text{S-HS}^{\text{act}}$ . Before or after pcDNA3 transfection, typically COS-7 conditioned Serum-Free Medium contained a low but detectable amount of  $\text{HS}^{\text{act}}$  conversion activity, whereas transfection by pCMV-3-OST elevated levels ~2,000-fold.

Second, to exclude the remote possibility that the expression of the mouse 3-OST-1 cDNA indirectly induces, rather than directly encodes,  $\text{HS}^{\text{act}}$  conversion activity, a Protein A/3-OST-1 fusion protein was analyzed. COS-7 cells were transiently transfected with pCMV-ProA3-OST, a pCMV-3-OST derivative in which the amino-terminal 26 residues of the mouse 3-OST-1 are replaced with a Protein A tag, and Protein A tagged mouse r3-OST-1 was extracted with IgG agarose beads from 155 ml of conditioned Serum-Free Medium, as described above. The affinity purification recovered undetectable and less than 0.5% of initial  $\text{HS}^{\text{act}}$  conversion activity from control pcDNA3 and pCMV-3-OST transfection samples, respectively, whereas ~7,000 units (10% recovery) were extracted from pCMV-ProA3-OST transfection samples. Thus, the mouse 3-OST-1 cDNA directly encodes  $\text{HS}^{\text{act}}$  conversion activity.

Third, the activities of cell-free synthesized mouse and human r3-OST-1 were examined. Synthetic capped mouse and human 3-OST-1 mRNAs were generated by *in vitro* transcription and then *in vitro* translated with reticulocyte lysate in the presence and absence of canine pancreatic microsomal membranes, as described above.  $\text{HS}^{\text{act}}$  conversion activity was undetectable in the control *in vitro* translation reactions which lacked mRNA template, with or without microsomal membranes. A low level  $\text{HS}^{\text{act}}$  conversion activity resulted from the addition of synthetic 3-OST-1 mRNA templates to translation reactions lacking microsomal membranes (mouse,  $0.86 \pm 0.028$  units/ $\mu\text{l}$ ,  $n = 3$ ; human,  $2.1 \pm 0.063$  units/ $\mu\text{l}$ ,  $n = 3$ ); however, ~15-fold greater levels occurred when microsomal membranes were included in translation reactions (mouse,  $14.3 \pm 0.27$  units/ $\mu\text{l}$ ,  $n = 3$ ; human,  $32.4 \pm 2.1$  units/ $\mu\text{l}$ ,  $n = 3$ ). The apparent activation of nascent r3-OST-1 by co-translational processing within microsomes may result from signal peptidase cleavage, *N*-linked glycosylation, and/or a facilitation of correct protein folding. The slightly greater production from the

human 3-OST-1 cDNA may reflect the more favorable context of the human initiation codon, or the reduced length of the human 5' untranslated region. Independent of the above considerations, the above data confirm that isolated mouse and human cDNAs encode HS<sup>act</sup> conversion activity.

5       Next, the biochemical specificity of the HS<sup>act</sup> conversion activity generated from each expression approach was examined by incubating crude or purified enzyme with [<sup>35</sup>S]PAPS and unlabeled HS<sup>inact</sup>, recovering radiolabeled GAG by DEAE chromatography and characterizing the resultant products. The HS<sup>act</sup> conversion activity of the wild-type mouse r3-OST-1 produced by transfecting COS-7 cells with  
10   pCMV-3-OST (1.35 x 10<sup>6</sup> units in 240 ml of conditioned Serum-Free Medium) was first purified away from potential contaminating sulfotransferase activities by heparin-AF Toyopearl chromatography followed by 3',5'-ADP-agarose chromatography, which yielded ~1 µg of protein containing 340,000 units (~20,000-fold purification with 25% overall recovery); whereas, the IgG agarose-purified Protein A tagged r3-OST-1  
15   and *in vitro* translation reactions of mouse and human 3-OST-1 mRNA templates were directly analyzed, as described above. About 0.5 - 1 x 10<sup>6</sup> cpm of product was generated with purified wild-type r3-OST-1, purified Protein A tagged r3-OST-1, and nonpurified *in vitro* translation reactions containing mouse and human r3-OST-1, respectively. Portions of each labeled product were incubated with purified  
20   heparitinase (0.5 units/ml) or chondroitinase ABC (0.5 units/ml) and HPLC-GPC analysis indicated that in all cases label was exclusively incorporated into HS. Portions of the labeled HS samples were also *N*-desulfated with nitrous acid at pH 1.5, and analyzed by P-2 polyacrylamide gel filtration to determine the amounts of liberated free [<sup>35</sup>S]sulfate, as described above. The results demonstrated no increased  
25   generation of free [<sup>35</sup>S]sulfate. Finally, portions of the labeled samples were AT affinity fractionated, which revealed that in each case ~40% of the <sup>35</sup>S-label was incorporated in HS<sup>act</sup> and approximately ~60% of the <sup>35</sup>S-label was incorporated in HS<sup>inact</sup>. The labeled HS<sup>act</sup> and HS<sup>inact</sup> generated by the wild-type purified r3-OST-1 were chemically cleaved to disaccharides with nitrous acid treatment, appropriate <sup>3</sup>H-  
30   labeled disaccharides standards were added, and the <sup>35</sup>S- and <sup>3</sup>H-labeled species were coresolved by RPIP-HPLC as outlined above. The results show that the <sup>35</sup>S-label coelutes with [<sup>3</sup>H]GlcA→AMN-3-*O*-SO<sub>3</sub> and [<sup>3</sup>H]GlcA→AMN-3,6-*O*-(SO<sub>3</sub>)<sub>2</sub>,

respectively. This approach also revealed that Protein A tagged r3-OST-1, and *in vitro* translation derived mouse and human r3-OST-1 generated  $^{35}\text{S}$ -HS which only contained  $^{35}\text{S}$ -labeled disaccharides that coeluted with  $[^3\text{H}]\text{GlcA}\rightarrow\text{AMN-3-O-SO}_3$  and  $[^3\text{H}]\text{GlcA}\rightarrow\text{AMN-3,6-O-(SO}_3)_2$ , respectively. It was previously shown that  $^{35}\text{S}$ -labeled  $\text{GlcA}\rightarrow\text{AMN-3,6-O-(SO}_3)_2$  generated by purified 3-OST-1 enzyme contains  $^{35}\text{S}$  solely in the 3-*O*- position (26). Thus, the expressed  $\text{HS}^{\text{act}}$  conversion activities exclusively catalyze the transfer of sulfate to the 3-*O*- position of glucosamine units in  $\text{HS}^{\text{act}}$  and  $\text{HS}^{\text{inact}}$ .

#### Northern Analysis of Rodent and Human 3-OST-1 Expression

Northern blot analysis reveals the presence of 3-OST-1 message in different kinds of endothelial cells as well as a mast cell line. Both cell types have previously been shown to form  $\text{HS}^{\text{act}}$  and anticoagulant heparin, respectively (6, 8, 55). Three size categories of rodent 3-OST-1 mRNA (about 1.7, 2.3, 3.3 kb) and a single size species of the human message (about 1.7 kb) were evident. As described above, the mouse forms arise from differential splicing within the 5' untranslated region. Similar size categories are also expressed by rat (RFPEC) endothelial cells, suggesting a similar mechanism of origin. The abundance of each category varies with each cell line, which suggests that a mechanism exists to regulate such differential splicing. The immortalized mouse mast cell line, C57.1, expresses high levels of the same three size categories, which suggests that expression of a single 3-OST-1 gene is required for the synthesis of both  $\text{HS}^{\text{act}}$  and anticoagulant heparin.

#### The 3-OST-1 Sequence Defines a Heparan Sulfotransferase Family

Extensive computer-aided data bank searching revealed the 3-OST-1 protein to be a previously unidentified protein; furthermore, the carboxy-terminal 250 residues exhibit a low homology (~30% similarity) to many previously identified sulfotransferases (which are typically ~300 residues in length) including chondroitin-, aryl-/phenol-, *N*-hydroxyarylamine-, alcohol-/hydroxysteroid-, flavonol-, and nodulation factor sulfotransferases. We also observed a slightly greater homology (~40% similarity) to a functionally unidentified open reading frame of 247 amino acids from *Aeromonas salmonicida* (GenBank accession number L37077). More importantly, the 3-OST-1 protein exhibits ~50% similarity with all previously identified forms of the heparan biosynthetic enzyme *N*-deacetylase/*N*-sulfotransferase

(NST). In particular, extensive homology exists across the entire 250-270 carboxy-terminal residues of these enzymes. Thus, it appears that a common sulfotransferase structure is shared by two distinct types of heparan biosynthetic enzyme. Given that NST is a bifunctional enzyme, the above observation suggests that NST enzymes possess sulfotransferase activity within a ~270 residue carboxy-terminal domain, whereas deacetylase activity would be contained within the remaining ~560 luminal residues. Interestingly, the region of consensus Lys<sup>302</sup>-Arg<sup>323</sup>, which encompasses the presumptive cysteine bridged peptide loop (described above), exhibits complete conservation for 12 of the 22 residues (including both cysteines) among all 3-OST-1 and NST species.

#### Identification and molecular cloning of 3-OST-2, 3-OST-3A, 3-OST-3B and 3-OST-4

The 3-OST-1 protein exhibits a COOH-terminal region of ~260 residues which was determined to be a sulfotransferase (ST) domain based on homology to all known sulfotransferases. The National Center for Biotechnology Information data bank of expressed sequence tags (ESTs) was searched with amino acid sequences of the ST domain from the human 3-OST-1 cDNA to reveal seven human cDNAs encoding three novel related species. The forms were subsequently designated as 3-OST-2 (I.M.A.G.E. Consortium (LLNL) CloneID c-20d10), 3-OST-3 (Clone ID 284542) and 3-OST-4 (Clone IDs HIBCX69, IB727, 166466, 23279, and c-3ie01). These EST clones were obtained from the TIGR/ATCC Special Collection, and the inserts were completely sequenced, revealing that all clones were of partial length.

To obtain full length clones, isoform specific probes were generated from the EST clones and used to screen  $\lambda$  TriplEx human cDNA libraries. 7 and 4 additional 3-OST-2 and 3-OST-4 cDNAs were isolated from a brain library, and 8 new 3-OST-3 cDNAs were recovered from a liver library. The cDNA inserts were completely sequenced, revealing the full length form for 3-OST-2 as well as 2 distinct full length forms for 3-OST-3 (3-OST-3A and 3-OST-3B). The additional 3-OST-4 clones were also of partial length.

#### 3-OST-2, 3-OST-3A, 3-OST-3B and 3-OST-4 Protein Structures and Activities

The 3-OST-2, 3-OST-3A, and 3-OST-3B proteins are 367, 406, and 390 amino acids in length, respectively. All three proteins conform to the architecture of a type-II integral membrane protein. These proteins and the partial length 3-OST-4

share a common (85% similarity) ST domain region of ~260 amino acid at their COOH-terminus. To characterize the encoded HS sulfotransferase activities, the 3-OST-2, 3-OST-3A, and 3-OST-3B cDNAs were individually expressed in COS-7 cells.

The analysis of transfected cell extracts demonstrated that each enzyme transfers sulfate specifically to the 3-O position of glucosamine residues within HS; however distinct specificities occur. 3-OST-2 preferentially sulfates regions containing GlcA 2S→GlcNS to generate GlcA 2S→GlcNS 3S; whereas both 3-OST-3A, and 3-OST-3B recognize regions with IdoA 2S→GlcNS to generate IdoA 2S→GlcNS 3S.

#### Expression Patterns Indicate Biological Function

The biologic function of these novel enzymes was elucidated by performing northern blot analysis. 3-OST-4 is exclusively expressed in the brain, whereas 3-OST-2 mRNA predominantly occurs in the brain with minor levels also found in heart, lung, skeletal muscle and placenta. 3-OST-3 forms occur in virtually all tissues but with barely detectable levels in brain, low levels in heart, lung, skeletal muscle and kidney, and extremely abundant expression in liver and placenta. Thus 3-OST-2 and 3-OST-4 appear to be the brain counterparts of 3-OST-3. The product of 3-OST-3 (IdoA 2S→GlcNS 3S) has previously been shown to be extremely abundant in HSPGs isolated from the glomerular basement membrane (GBM) of the kidney. These HSPGs are critical to regulating the permselectivity of the GBM. This function occurs through interactions with extracellular matrix components that regulate the pore size of the matrix. Given that the liver, placenta, and kidney glomerulus are all responsible for the filtration of macromolecular components from blood and all exhibit high 3-OST-3 expression, it appears that 3-OST-3 serves a common function in each situation: to regulate macromolecular permeability. In this functional regard, the high brain expression of 3-OST-2 and 3-OST-4 correlates with the major molecular permeability barrier of the central nervous system, the blood brain barrier.

#### Therapeutic Utilities

The 3-OST heparan biosynthetic enzymes may be generated by recombinant expression of the isolated cDNAs to generate novel glycosaminoglycan drugs of specific structure through an *in vitro* biochemical synthesis approach. Specifically, 3-

OST-1 may be used to generate anticoagulant pentasaccharides, which may be administered subcutaneously to treat thrombotic disorders such as deep vein thrombosis and pulmonary embolism. The 3-OST-1 enzyme may also be used to generate an orally absorbable form of pentasaccharide from an appropriate carbohydrate substrate linked to a hydrophobic group. In an analogous fashion, specific glycosaminoglycan products may be generated from 3-OST-2, 3-OST-3 and 3-OST-4, which may be used as therapeutics to alter macromolecular permeability of various vascular beds. Drugs which reduce capillary permeability may, at the very least, be used to treat (1) microproteinuria and macroproteinuria of renal diseases including diabetic nephropathy and the various forms of glomerulonephritis; (2) neoplastic growths by limiting nutrient supply to tumors; and (3) inflammatory diseases where macromolecular constituents of the plasma are required for initiating and maintaining a localized inflammation. Conversely, drugs which enhance capillary permeability may be used (1) as an adjunctive treatment to facilitate pharmacological access to vascular beds, which exhibit highly selective drug entry, such as the blood brain barrier and the placental barrier; and (2) to enhance nutrient supply to under-perfused tissues such as the myocardium after an infarct.

Specific heparan sulfate structures regulate additional biologic processes by interacting with numerous protein effector molecules including growth and differentiation factors (e.g., FGF family members, HB-EGF, HGF/SF, interferon  $\gamma$ , PDGF, SDGF, and VEGF/VPF), chemokines (e.g., MIP-1 $\beta$ , RANTES, and GRO), receptors (e.g., TGF- $\beta$  receptors), mast cell proteases, protease inhibitors (e.g., AT, heparin cofactor II, leuserpin, plasminogen activator inhibitor-1, protease nexins), degradative enzymes (e.g., elastase, acetylcholinesterase, extracellular superoxide dismutase, thrombin, tissue plasminogen activator, lipoprotein lipase, hepatic and pancreatic triglyceride lipase, and cholesterol esterase), apolipoproteins (e.g., apoB and apoE), matrix components (e.g., fibronectin, wnt-1, interstitial collagens, laminin, pleiotropin, tenascin, thrombospondin, and vitronectin) viral coat proteins (e.g., gC and gB of HSV types I and II, gC-II of CMV, and gp120 of HIV), nuclear proteins (e.g., c-fos, c-jun, RNA and DNA polymerases, and steroid receptors), cellular adhesion molecules (e.g., L-selectin, P-selectin, PECAM-1, and N-CAM) and other molecules (e.g., HB-GAM/pleiothrophin, amphoterin, and PF4).

Using routine methods (e.g., site-directed mutagenesis) the available 3-OST cDNAs may be selectively mutated to alter substrate recognition properties so as to produce enzymes that generate novel glycosaminoglycan structures which modulate the biologic processes regulated by the above effector molecules. Thus, novel drugs may also be biochemically synthesized from recombinantly expressed mutated enzymes. Such substances may serve to (1) enhance growth or regeneration of specific cell types such as the endothelial cells of the heart after infarction, or neurons in neurodegenerative diseases; (2) suppress undesirable cell growth in conditions such as cancer (either directly by acting on the cancers cells or indirectly by preventing endothelial cells from neovascularizing the tumor), atherosclerosis (by preventing smooth muscle cell growth), and inflammatory diseases characterized by cellular proliferation; (3) prevent metastasis of tumors by modulating cell/matrix interactions; (4) reduce the destructive side effects of inflammatory reactions by inhibiting degradative enzymes or by activating inhibitory molecules (e.g. protease inhibitors) which may be directly or indirectly protective by limiting extravasation of lymphocytes; (5) modulate serum lipid levels by enhancing or reducing the cellular or tissue uptake or degradation of specific lipoprotein classes; (6) treat viral infections by preventing viral entry into cells; and (7) facilitate axon regeneration subsequent to nerve severing.

Bacterial expression of 3-OST1. The human and mouse 3-OST-1 proteins have been expressed as active, soluble protein in *E. coli*. This has been achieved using the pET system from NOVEGEN (Madison, WI). The human and mouse 3-OST-1 cDNA's were PCR amplified with pfu DNA polymerase and purified cloned plasmids as template. The primers that were used were designed to amplify a cDNA fragment starting, in frame, after the native signal sequence and including the native translational termination codon. Additionally, the PCR primers were designed to include restriction sites that would facilitate cloning into the vectors described below in the correct transcriptional/translational reading frames. 3-OST-1 was cloned into vectors pET12a, 15B and 28a according to the manufacturer's instructions. This places the 3-OST-1 cDNA downstream of a powerful, inducible T7 transcription site and includes an efficient Shine-Dalgarno sequence at the appropriate distance from the initiator methionine of the construct.



Good yields of active protein result from IPTG induction at room temperature.

The specific activity appears to be less than purified, or Baculovirus/sf9 produced material. The exact magnitude of the diminution of activity is unclear at this time; however, it may be 10-1000 fold. The presently preferred purification scheme is: (1)

- 5 Induction at 22 °C. (2) Sonication of bacteria, centrifugation to remove inclusion bodies and cell debris, purification of crude bacterial sonicate on heparin sepharose as described elsewhere. (3) PAP column chromatography. (4) Gel permeation chromatography. Step (4) is only needed for obtaining monomeric, pure 3-OST-1, and not for active protein preparation.

095736-0440  
004240" 2924560

## REFERENCES

1. Morawitz, P. (1958) *The Chemistry of Blood Coagulation.*, Charles C Thomas, Springfield, IL
2. Rosenberg, R. D., and Damus, P. S. (1973) *J. Biol. Chem.* **248**, 6490-6505
- 5 3. Damus, P. S., Hicks, M., and Rosenberg, R. D. (1973) *Nature* **246**, 355-357
4. Marcum, J. A., McKenney, J. B., Galli, S. J., Jackman, R. W., and Rosenberg, R. D. (1986) *Am. J. Physiol.* **250**, H879-H888
5. Marcum, J. A., McKenney, J. B., and Rosenberg, R. D. (1984) *Journal of Clinical Investigation* **74**, 341-350
- 10 6. Marcum, J. A., Atha, D. H., Fritze, L. M. S., Nawroth, P., Stern, D., and Rosenberg, R. D. (1986) *J. Biol. Chem.* **261**, 7507-7517
7. Marcum, J. A., and Rosenberg, R. D. (1985) *Biochem. Biophys. Res. Commun.* **126**, 365-372
8. Kojima, T., Leone, C. W., Marchildon, G. A., Marcum, J. A., and Rosenberg, R. D. (1992) *J. Biol. Chem.* **267**, 4859-4869
- 15 9. Lindahl, U., and Kjellén, L. (1987) in *The Biology of the Extracellular Matrix Proteoglycans* (Wight, T. N., and Mecham, R., eds), pp. 59-104, Academic, New York
10. Atha, D. H., Stevens, A. W., Rimon, A., and Rosenberg, R. D. (1984) *Biochemistry* **23**, 5801-5812
- 20 11. Atha, D. H., Lormeau, J. C., Petitou, M., Rosenberg, R. D., and Choay, J. (1985) *Biochemistry* **24**, 6723-6729
12. Atha, D. H., Lormeau, J. C., Petitou, M., Rosenberg, R. D., and Choay, J. (1987) *Biochemistry* **26**, 6454-6461
- 25 13. Choay, J., Petitou, M., Lormeau, J. C., Sinaÿ, P., Casu, B., and Gatti, G. (1983) *Biochem. Biophys. Res. Commun.* **116**, 492-499
14. Lindahl, U., Bäckström, G., Thunberg, L., and Leder, I. G. (1980) *Proc. Nat. Acad. Sci. U. S. A.* **77**, 6551-6555
15. Lindahl, U., Bäckström, G., and Thunberg, L. (1983) *J. Biol. Chem.* **258**, 9826-9830
- 30 16. Rosenberg, R. D., and Lam, L. (1979) *Proc. Nat. Acad. Sci. U. S. A.* **76**, 1218-1222

17. Rosenberg, R. D., Armand, G., and Lam, L. (1978) *Proc. Nat. Acad. Sci. U. S. A.* **75**, 3065-3069
18. Oosta, G. M., Gardner, W. T., Beeler, D. L., and Rosenberg, R. D. (1981) *Proc. Nat. Acad. Sci. U. S. A.* **78**, 829-833
- 5 19. Stone, A. L., Beeler, D. L., Oosta, G. M., and Rosenberg, R. D. (1982) *Proc. Nat. Acad. Sci. U. S. A.* **79**, 7190-7194
20. Lind, T., Lindahl, U., and Lidholt, K. (1993) *J. Biol. Chem.* **268**, 20706-20708
21. Brandan, E., and Hirschberg, C. B. (1988) *J. Biol. Chem.* **263**, 2417-2422
22. Pettersson, I., Kusche, M., Unger, E., Wlad, H., Nylund, L., Lindahl, U., and
- 10 Kjellén, L. (1991) *J. Biol. Chem.* **266**, 8044-8049
23. Campbell, P., Hannesson, H. H., Sandbäck, D., Rodén, L., Lindhal, U., and Li, J.-P. (1994) *J. Biol. Chem.* **269**, 26953-26958
24. Kobayashi, M., Habuchi, H., Habuchi, O., Saito, M., and Kimata, K. (1996) *J. Biol. Chem.* **271**, 7645-7653
- 15 25. Habuchi, H., Habuchi, O., and Kimata, K. (1995) *J. Biol. Chem.* **270**, 4172-4179
26. Liu, J., Shworak, N. W., Fritze, L. M. S., Edelberg, J. M., and Rosenberg, R. D. (1996) *J. Biol. Chem.* **271**, 27072-27082
27. Orellana, A., Hirschberg, C. B., Wei, Z., Swiedler, S. J., and Ishihara, M. (1994) *J. Biol. Chem.* **269**, 2270-2276
- 20 28. Hashimoto, Y., Orellana, A., Gil, G., and Hirschberg, C. B. (1992) *J. Biol. Chem.* **267**, 15744-15750
29. Eriksson, I., Sandbäck, D., Ek, B., Lindahl, U., and Kjellén, L. (1994) *J. Biol. Chem.* **269**, 10438-10443
30. Ishihara, M., Guo, Y., Wei, Z., Yang, Z., Swiedler, S. J., Orellana, A., and
- 25 Hirschberg, C. B. (1993) *J. Biol. Chem.* **268**, 20091-20095
31. Cheung, W.-F., Eriksson, I., Kusche-Gullberg, M., Lindahl, U., and Kjellén, L. (1996) *Biochemistry* **35**, 5250-5256
32. Casu, B. (1985) *Adv. Carbohydr. Chem. Biochem.* **43**, 51-134
33. Shworak, N. W., Shirakawa, M., Collic-Jouault, S., Liu, J., Mulligan, R. C.,
- 30 Birinyi, L. K., and Rosenberg, R. D. (1994) *J. Biol. Chem.* **269**, 24941-24952
34. Collic-Jouault, S., Shworak, N. W., Liu, J., de Agostini, A. I., and Rosenberg, R. D. (1994) *J. Biol. Chem.* **269**, 24953-24958

35. Shworak, N. W., Fritze, L. M. S., Liu, J., Butler, L. D., and Rosenberg, R. D. (1996) *J. Biol. Chem.* **271**, 27063-27071
36. Lam, L. H., Silbert, J. E., and Rosenberg, R. D. (1976) *Biochem. Biophys. Res. Commun.* **69**, 570-577
- 5 37. Linhardt, R. J., Wang, H., Loganathan, D., and Bae, J. (1992) *J. Biol. Chem.* **267**, 2380-2387
38. Kusche, M., Torri, G., Casu, B., and Lindahl, U. (1990) *J. Biol. Chem.* **265**, 7292-7300
39. Montgomery, R., Lidholt, K., Flay, N., Liang, J., Vertel, B., Lindahl, U., and
- 10 Esko, J. (1992) *Proc. Nat. Acad. Sci. U. S. A.* **89**, 11327-11331
40. Razi, N., and Lindahl, U. (1995) *J. Biol. Chem.* **270**, 11267-11275
41. de Agostini, A. L., Lau, H. K., Leone, C., Youssofian, H., and Rosenberg, R. D. (1990) *Proc. Nat. Acad. Sci. U. S. A.* **87**, 9784-9788
42. Shworak, N. W., Shirakawa, M., Mulligan, R. C., and Rosenberg, R. D. (1994) *J.*
- 15 *Biol. Chem.* **269**, 21204-21214
43. Fernandez, J., DeMott, M., Atherton, D., and Mische, S. M. (1992) *Anal. Biochem.* **201**, 255-264
44. Stone, K. L., Elliott, J. I., Peterson, G., McMurray, W., and Williams, K. R. (1990) *Methods Enzymol.* **193**, 389-412
- 20 45. Favaloro, J., Treisman, R., and Kamen, R. (1980) *Methods Enzymol.* **65**, 718-749
46. Hecker, K. H., and Roux, K. H. (1996) *BioTechniques* **20**, 478-485
47. Don, R. H., Cox, P. T., Wainwright, B. J., Baker, K., and Mattick, J. S. (1991) *Nucleic Acids Res.* **19**, 4008
48. Lennon, G., Auffray, C., Polymeropoulo, M., and Soares, M. B. (1996) *Genomics*
- 25 **33**, 151-152
49. Wei, Z., Swiedler, S. J., Ishihara, M., Orellana, A., and Hirschberg, C. B. (1993) *Proc. Nat. Acad. Sci. U. S. A.* **90**, 3885-3888
50. Kojima, T., Shworak, N. W., and Rosenberg, R. D. (1992) *J. Biol. Chem.* **267**, 4870-4877
- 30 51. Chomczynski, P., and Sacchi, M. (1987) *Anal. Biochem.* **162**, 156-159
52. Young, J. D.-E., Liu, C.-C., Butler, G., Cohn, Z. A., and Galli, S. J. (1987) *Proc. Nat. Acad. Sci. U. S. A.* **84**, 9175-9179

53. Kozak. (1989) *J. Cell Biol.* **108**, 229-241
54. von Heijne, G. (1986) *Nucleic Acids Res.* **14**, 4683-4690
55. Robinson, H. C., Horner, A. A., Höök, M., Ögren, S., and Lindahl, U. (1978) *J. Biol. Chem.* **253**, 6687-6693
- 5 56. Andres, D. A., Rhodes, J. D., Meisel, R. L., and Dixon, J. E. (1991) *J. Biol. Chem.* **266**, 14277-14282
57. Munro, S., and Pelham, H. R. (1987) *Cell* **48**, 899-907
58. Weis, K., Griffiths, G., and Lamond, A. I. (1994) *J. Biol. Chem.* **269**, 19142-19150
- 10 59. Scherer, P. E., Lederkremer, G. Z., Williams, S., Fogliano, M., Baldini, G., and Lodish, H. F. (1996) *J. Cell Biol.* **133**, 257-268
60. Hong, W., and Tang, B. L. (1993) *BioEssays* **15**, 231-238
61. Machamer, C. E. (1991) *Trends Cell Biol.* **1**, 141-144
62. Weinshilboum, R. M., Otterness, D. M., Aksoy, I. A., Wood, T. C., Her, C., and
- 15 Raftogianis, R. B. (1997) *FASEB J.* **11**, 3-14
63. Walker, J. E., Saraste, M., Runswick, M. J., and Gay, N. J. (1982) *EMBO J.* **1**, 945-951
64. Saraste, M., Sibbald, P. R., and Wittinghofen, A. (1990) *Trends Biochem. Sci.* **15**, 430-434
- 20 65. Zheng, Y., Bergold, A., and Duffel, M. W. (1994) *J. Biol. Chem.* **269**, 30313-30319
66. Habuchi, O., Tsuzuki, M., Takeuchi, I., Hara, M., Matsui, Y., and Ashikari, S. (1991) *Biochim. Biophys. Acta* **1133**, 9-16
67. Nishinaga, M., Ozawa, T., and Shimada, K. (1993) *J. Clin. Invest.* **92**, 1381-1386
- 25 68. Wilson, R., et al. (1994) *Nature* **368**, 32-38
69. Hogan, et al. (1986) Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York
70. Shworak et al. (1997) *J. Biol. Chem.* **272**, 28008-28019

## CLAIMS

- 1 ~~1.~~ An isolated nucleic acid encoding at least a functional fragment of a 3-OST  
2 protein.
- 1 2. An isolated nucleic acid as in claim 1 wherein said nucleic acid encodes a 3-  
2 OST protein comprising a mature 3-OST-1 protein selected from the group consisting  
3 of mature murine 3-OST-1 and mature human 3-OST-1.
- 1 3. An isolated nucleic acid as in claim 1 wherein said nucleic acid encodes a 3-  
2 OST protein comprising a protein selected from the group consisting of 3-OST-1, 3-  
3 OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4, and ce3-OST.
- 1 4. An isolated nucleic acid as in claim 1 wherein said nucleic acid encodes a 3-O-  
2 sulfotransferase domain of a 3-OST protein selected from the group consisting of 3-  
3 OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4, ce3-OST.
- 1 5. An isolated nucleic acid as in claim 1 wherein said nucleic acid comprises a  
2 nucleotide sequence selected from nucleotide sequences within:  
3 (a) SEQ ID NO: 1;  
4 (b) SEQ ID NO: 3;  
5 (c) SEQ ID NO: 5;  
6 (d) SEQ ID NO: 7;  
7 (e) SEQ ID NO: 9;  
8 (f) SEQ ID NO: 11;  
9 (g) a sequence having at least 60% nucleotide sequence identity with at least  
10 one of (a)-(f) and encoding a functional fragment having sequence-specific HS  
11 binding affinity or 3-O-sulfotransferase activity; and  
12 (h) a sequence differing from a sequence of (a)-(g) only by the substitution of  
13 synonymous codons.
- 1 6. An isolated nucleic acid as in claim 1 wherein said nucleic acid comprises a  
2 nucleotide sequence encoding a polypeptide selected from the group consisting of:  
3 (a) residues 21-52, 260-269, 250-276, 53-311, or 21-307 of SEQ ID NO: 2;  
4 (b) residues 21-48, 256-265, 246-272, 49-307, or 21-303 of SEQ ID NO: 4;  
5 (c) residues 42-109, 313-325, 303-332, or 110-367 of SEQ ID NO: 6;  
6 (d) residues 44-147, 351-363, 341-370, or 148-406 of SEQ ID NO: 8;

(e) residues 66-132, 336-348, 326-355, or 133-390 of SEQ ID NO: 10;

(f) residues 396-408, 386-4150, or 207-456 of SEQ ID NO: 12;

(g) residues 240-250, 230-257, 23-291 of SEQ ID NO: 15;

(h) a sequence having at least 60% amino acid sequence similarity with at least one of (a)-(g) and encoding a functional fragment having sequence-specific HS binding affinity or 3-O-sulfotransferase activity; and

(i) a sequence comprising a chimera of at least two of sequences (a)-(h).

7. An isolated nucleic acid comprising at least 16 consecutive nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, and SEQ ID NO: 11.

8. A host cell transformed with a nucleic acid of any one of claims 1-7, or a descendant thereof.

9. A host cell as in claim 8 wherein said host cell is selected from the group consisting of bacterial cells, yeast cells, and insect cells.

10. A host cell as in claim 8 wherein said host cell is selected from the group consisting of somatic cells, fetal cells, embryonic stem cells, zygotes, gametes, germ line cells, and transgenic animal cells.

11. A host cell as in claim 8 wherein said cell is a mammalian cell.

12. A host cell as in claim 11 wherein said cell is selected from the group consisting of: COS-7 cells, CHO, murine primary cardiac microvascular endothelial cells (CME), murine mast cell line C57.1, human primary endothelial cells of umbilical vein (HUVEC), F9 embryonal carcinoma cells, rat fat pad endothelial cells (RFPEC), L cells, and cells derived from the transgenic animals of the invention.

13. A substantially pure protein preparation comprising at least a functional fragment of a 3-OST protein.

14. A substantially pure protein preparation as in claim 13 wherein said 3-OST protein is selected from the group consisting of mature murine 3-OST-1 and mature human 3-OST-1.





(h) a sequence having at least 60% amino acid sequence similarity with at least one of (a)-(g) and encoding a functional fragment having sequence-specific HS binding affinity or 3-O-sulfotransferase activity; and

(i) a sequence comprising a chimera of at least two of sequences (a)-(h).

~~19.~~ A method of 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides comprising:

contacting said preparation with at least a 3-O-sulfotransferase domain of a 3-OST protein in the presence of a sulfate donor under conditions which permit sulfation of said residues;

wherein, said 3-OST protein is selected from the group consisting of 3-OST-1, 3-OST-2, 3-OST-3A, 3-OST-3B, 3-OST-4, ce3-OST, and conservative substitution variants or chimeras thereof.

~~20.~~ A method of 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides, wherein said polysaccharides include a polysaccharide sequence of GlcA→GlcNS ±6S comprising:

contacting said preparation with a 3-OST-1 protein in the presence of a sulfate donor under conditions which permit said 3-OST-1 to convert said GlcA→GlcNS ±6S sequence to GlcA→GlcNS 3S ±6S.

wherein the 3-OST-1 protein is selected from the group consisting of murine 3-OST-1, human 3-OST-1, mature murine 3-OST-1, mature human 3-OST-1, a functional fragment of a 3-OST-1 having 3-O-sulfotransferase activity, a conservative substitution variant of 3-OST-1 having 3-O-sulfotransferase activity, and a chimeric 3-OST-1 having 3-O-sulfotransferase activity.

21. A method as in claim 20, wherein said GlcA→GlcNS ±6S polysaccharide sequence comprises a part of a polysaccharide sequence selected from the group consisting of:

(a) GlcA→GlcNS ±6S→IdoA 2S→ GlcNS ±6S;

(b) IdoA→GlcNAc 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S;

(c) IdoA→GlcNS 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S;

(d) IdoA→GlcNAc→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S;

(e) IdoA→GlcNS→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS 6S;

(f) IdoA→GlcNAc 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS;

(g) IdoA→GlcNS 6S→GlcA→GlcNS ±6S→IdoA 2S→ GlcNS;

~~22.~~ A method of 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides, wherein said polysaccharides include a polysaccharide sequence of GlcA 2S→GlcNS comprising:

contacting said preparation with a 3-OST-2 protein in the presence of a sulfate donor under conditions which permit said 3-OST-2 to convert said GlcA 2S→GlcNS sequence to GlcA 2S→GlcNS 3S.

wherein the 3-OST-2 protein is selected from the group consisting of 3-OST-2, a functional fragment of a 3-OST-2 having 3-O-sulfotransferase activity, a conservative substitution variant of 3-OST-2 having 3-O-sulfotransferase activity, and a chimeric 3-OST-2 having 3-O-sulfotransferase activity.

23. A method as in claim 22, wherein said GlcA 2S→GlcNS polysaccharide sequence comprises a part of a GlcNS→GlcA 2S→GlcNS polysaccharide sequence.

~~24.~~ A method of 3-O-sulfating saccharide residues within a preparation of glycosaminoglycan or proteoglycan polysaccharides, wherein said polysaccharides include a polysaccharide sequence of IdoA 2S→GlcNS comprising:

contacting said preparation with a 3-OST-3 protein in the presence of a sulfate donor under conditions which permit said 3-OST-3 to convert said IdoA 2S→GlcNS sequence to IdoA 2S→GlcNS 3S.

wherein the 3-OST-3 protein is selected from the group consisting of 3-OST-3A, 3-OST-3B, a functional fragment of a 3-OST-3 having 3-O-sulfotransferase activity, a conservative substitution variant of 3-OST-3 having 3-O-sulfotransferase activity, and a chimeric 3-OST-3 having 3-O-sulfotransferase activity.

25. A method as in claim 24, wherein said IdoA 2S→GlcNS polysaccharide sequence comprises a part of a GlcNS→IdoA 2S→GlcNS polysaccharide sequence.

~~26.~~ A method for enriching the AT-binding fraction in a preparation of heparan sulfates, wherein said preparation includes a polysaccharide sequence of GlcA→GlcNS ±6S comprising:

4 contacting said preparation with 3-OST-1 protein in the presence of a sulfate  
5 donor under conditions which permit said 3-OST-1 to convert said GlcA→GlcNS ±6S  
6 sequence to GlcA→GlcNS 3S ±6S, thereby increasing the fraction of AT-binding  
7 heparan sulfates.

1 ~~27.~~ A method for converting HS<sup>act</sup> precursor to HS<sup>act</sup> in a preparation of heparan  
2 sulfates, wherein said preparation includes HS<sup>act</sup> precursor polysaccharides including  
3 a sequence of GlcA→GlcNS ±6S comprising:

4 contacting said preparation with 3-OST-1 protein in the presence of a sulfate  
5 donor under conditions which permit said 3-OST-1 to convert said GlcA→GlcNS ±6S  
6 sequence to GlcA→GlcNS 3S ±6S, thereby converting HS<sup>act</sup> precursor to HS<sup>act</sup>.

1 28. A method as in any one of claims 19-28 wherein said sulfate donor is PAPS.

1 ~~29.~~ A non-human animal model, wherein a genome of said animal, or an ancestor  
2 thereof, wherein said recombinant construct has introduced a modification into said  
3 genome, said modification selected from the group consisting of insertion of a nucleic  
4 acid encoding at least a functional fragment of a conspecific wild type 3-OST protein,  
5 insertion of a nucleic acid encoding at least a functional fragment of a transpecific  
6 allelic variant of the 3-OST sequences, insertion of nucleic acid encoding at least a  
7 functional fragment of an allelic variant of 3-OST sequence, inactivation of an  
8 endogenous 3-OST gene, and insertion by homologous recombination of a reporter  
9 gene coupled to 3-OST transcriptional elements.

1 30. An animal as in claim 29 wherein said modification is insertion of nucleic acid  
2 encoding at least a functional fragment of wild type 3-OST selecting from the  
3 sequence consisting of the SPLAG-domain, the cysteine-binding peptide loop, and the  
4 ~260 residue ST domain.

1 31. An animal as in claim 29 wherein said animal is selected from the group  
2 consisting of rats, mice, hamsters, guinea pigs, rabbit, dogs, cats, goats, sheep, pigs,  
3 and non-human primates.

1 32. An animal as in claim 29 wherein said animal is an invertebrate.

1 ~~33.~~ A method of producing antibodies which selectively bind to a 3-OST protein  
2 comprising the steps of

3 administering an immunogenically effective amount of a 3-OST epitope to an  
4 animal;  
5 allowing said animal to produce antibodies to said epitope; and  
6 obtaining said antibodies from said animal or from a cell culture derived  
7 therefrom.

1 34 A substantially pure preparation of antibody which selectively binds to an  
2 epitope of a 3-OST protein.

1 35. A substantially pure preparation of an antibody as claimed in 34 wherein said  
2 antibody selectively binds to at least a fragment of 3-OST.

1 36. A cell line producing an antibody of any one of the claims 34.

1 37. A method for identifying compounds which can modulate the expression of a  
2 3-OST gene comprising steps of  
3 providing a cell expressing a nucleic acid under the control of a 3-OST  
4 regulatory sequence;

5       contacting said cell with at least one candidate compound; and  
6       assaying for a change in the in the expression of said nucleic acid.

1 38. The method of claim 37, wherein said nucleic acid comprises a marker gene  
2 and a 3-OST gene

1 39. The method of claim 37, wherein said said assaying step comprises detecting a  
2 change in 3-OST mRNA level.

1 40. The method of claim 37, wherein said assaying step comprises detecting a  
2 change in 3-OST protein encoded by said nucleic acid.

1 41. A method of determining partial sequence information for complex  
2 polysaccharides comprising the steps of:

3           contacting a first sample of polysaccharide with at least one ligand which  
4           binds polysaccharides in a sequence specific manner;

5           contacting the resulting polysaccharide-ligand complex with at least one agent  
6   that modifies complex polysaccharides;

7 contacting a second sample of polysaccharide with the same modifying agent;

comparing said first and second samples for ligand-specific inhibition of modifications caused by said modifying agent.

42. The method of claim 41, wherein said complex polysaccharide is a glycosaminoglycan.

43. The method of claim 41, wherein said ligand is catalytically inactive.

44. The method of claim 41, wherein said ligand is an inactive 3-OST.

45. The method of claim 41, wherein said agent that modifies polysaccharides is selected from the group consisting of epimerases, lyases, sulfotransferases, N-actyltransferases, N-deacetylases, epimerases.

46. The method of claim 45, wherein said modifying agent is a sequence specific degrading agent.

47. The method of claim 45, wherein said modifying agent is a non-sequence specific degrading agent.

48. The method of claims 46, wherein said degrading agent is a lyase.

49. The method of claim 47, wherein said non-sequence specific degrading agent is nitrous acid.

50. The method of claim 45, further comprising affinity purifying said modified first and second samples.

51. The method of claim 45, wherein the step of comparing includes a comparison of size profiles.

52. A method of determining partial sequence information for complex polysaccharides comprising the steps of:

contacting a first sample of complex polysaccharides with a 3-OST protein in the presence of a sulfate donor under conditions which permit sulfation by said 3-OST;

contacting said first sample and a second sample with at least one enzyme which cleaves polysaccharides in a sequence-specific manner;

determining the size profiles of the resulting fragments.

53. The method of claim 52, wherein the determining the size profile step further comprises the step of comparing said first sample to a second sample cleaved by the same enzymes.

1 54. The method of claim 52, wherein said enzymes which degrade polysaccharides  
2 in a sequence specific manner are selected from the group consisting of  
3 polysaccharide lyases, heparinase I, heparinase II, and heparinase III

1 55. A method of determining partial sequence information for a sample containing  
2 complex polysaccharides comprising the steps of:  
3 contacting said sample of polysaccharide with a 3-OST protein which lacks  
4 enzymatic function with a under conditions which permit said 3-OST protein to bind  
5 to said polysaccharide in a sequence specific manner;  
6 applying said sample to an affinity column;  
7 applying degrading agents to said column;  
8 analyzing the resulting degradation products.

1 56. The method of claim 55, further comprising repeating the steps applying  
2 degrading agents and analyzing using a series of different sequence specific  
3 polysaccharide cleavage enzymes.

1 57. An isolated nucleic acid comprising a genetic regulatory sequences of a 3-OST  
2 operably joined to a marker gene.

1 58. A host cell transformed with the isolated nucleic acid of claim 57, or a  
2 descendent thereof.

1 59. A method of identifying compounds capable of modulating the expression of a  
2 3-OST comprising contacting candidate compound seith the transformed host cell of  
3 claim 58 and assaying for changes in expression of said marker.

1 60. A method as in claim 59, wherein said regulatory sequences comprise the 5'  
2 untranslated region of SEQ. ID NO: 16.

# HEPARAN SULFATE D-GLUCOSAMINYL 3-O-SULFOTRANSFERASES, AND USES THEREFOR

## Abstract of the Disclosure

Disclosed are novel isolated nucleic acids and substantially pure protein preparations for naturally occurring and synthetic or chimeric heparan sulfate D-glucosaminy 3-O-sulfo-transferases (3-OSTs). Also disclosed are uses for these genes and proteins, including uses for the modification and sequencing of

5 glycosaminoglycans.

BRESNAHA5473/91.984810

004240" 2326560



m3-OST-1	MTLLLLGAVL	LVAQPQLVHS	HPAAPGPGLK	QQELLRKVII	40
h3-OST-1	MAALLLGAVL	LVAQPQLVPS	RPA---ELG	QQELLRKAGT	36
m3-OST-1	LPEDTGEGTA	SNGSTQQLPQ	TIIIGVRKGG	TRALLEMLSL	80
h3-OST-1	LQDDVRDGVA	PNGSAQQLPQ	TIIIGVRKGG	TRALLEMLSL	76
m3-OST-1	HPDVAAAENE	VHFFDWEEHY	SQGLGWYLTQ	MPFSSPHQLT	120
h3-OST-1	HPDVAAAENE	VHFFDWEEHY	SHGLGWYLSQ	MPFSWPHQLT	116
m3-OST-1	VEKTPAYFTS	PKVPERIHSM	NPTIRLLLIL	RDPSEVLSD	160
h3-OST-1	VEKTPAYFTS	PKVPERVYSM	NPSIRLLLIL	RDPSEVLSD	156
m3-OST-1	YTQVLYNHLQ	KHKPYPIED	LLMRDGRNL	DYKALNRSly	200
h3-OST-1	YTQVFYNHMQ	KHKPYPSIEE	FLVRDGRNLV	DYKALNRSly	196
m3-OST-1	HAHMLNWLRF	FPLGHIHIVD	GDRLIRDPFP	EIQKVERFLK	240
h3-OST-1	HVHMQNWLRF	FPLRHIHIVD	GDRLIRDPFP	EIQKVERFLK	236
m3-OST-1	LSPQINASNF	YFNKTKGFYC	LRDSGKDRCL	HESKGRAHPQ	280
h3-OST-1	LSPQINASNF	YFNKTKGFYC	LRDSGRDRCL	HESKGRAHPQ	276
m3-OST-1	VDPKLLDKLH	EYFHEPNKKF	FKLVGRTFDW	H	311
h3-OST-1	VDPKLLNKLH	EYFHEPNKKF	FELVGRTFDW	H	307

Figure 1

599	KTCDFPKLLIIGPÖKTGTALYLFLGMHPDLSNYPSSSETEEEIOFFEN	GHNYHKGIDWMEFFPIPSNTTSDFYEKS	NST-1
598	KTCDRLPKFLIVGÖKTGTALHFFLSLHPAVTSSFPSPSTFEEIOFFEN	SPNYHKGIDWYMDFFVPASNASTDFLEKS	NST-2
23	TSKRFPDAIIVGVKKSSTRALLEFLRVNPLI	KAPPEVHEFDKNEN	ce3-OST
193	GEKK LPÖALIIGVKKGGTRALLEAIRVHPDV	RAVGEPHEFDNRN	3-OST-4
148	GS KÖLPÖALIIGVKKGGTRALLEFLRVHPDV	RAVGAEPHEFDRS	3-OST-3A
110	GT KRLPÖALIIVGVKKGGTRAVLEFIRVHPDV	RALGTEPHEFDRN	3-OST-2
49	GSAÖÖLPÖTIIIGVRKGGTRALLEMLSLHPDV	AAAEVHEFFDWEHSHGLGWTLSÖMPFSWP	3-OST-1
678	ANYEDSEVAPRRRAALLPKAKVLTILINPADRAYSWYÖHÖRAHDDPVALKYTFHEVITAGSDA	SSKLRALÖ	NST-1
677	ATYEDSEVAPRRGAALLPRAKITIVLTINPADRAYSWYÖHÖRAHGDPVALNTFYÖVISASSÖT	PLALRSLÖ	NST-2
93	PAYEHSKMAPERIKSLINENTKIIIVRDPVTRAISDYTÖSSSKRKRVGLM	PSEETMAVGNCANWLRNTCTTKTRGVNAG	ce3-OST
262	PSYFVTNEAPKRIHSMAKDIKLIVVRNPNVTRAISDYTÖT	LSKRPEIPTEEVLAFKNRT	3-OST-4
217	PSYFVTREAPARISAMSKDTKLIVVRDPVTRAISDYTÖT	LSKRDPDIPTEESLTFKNRT	3-OST-3A
179	PSYFVTOEAPRIENMSRDTKLIVVRNPNVTRAISDYTÖT	LSKRPDIPTEEGLSFRNRT	3-OST-2
121	PAYFTSPKVPERVYSMNPSIRLLILRDPSEKRVLSDYTÖVFYNNHMQKHKPYPSIEEFLVRD	LSKRPDIPTEEGLSFRNRT	3-OST-1
749	NRCILVPGWYATHIERWLSAYHANÖILVLGDKLIRTEPAKMDVÖKFLGVTNTIDYHKTIAFDPKKGFWC	ÖLLEGKGT	NST-1
748	NRCILVPGYXSTHÖRWLTYPPSGÖLLIVDGÖELRTNPAASMESIÖKFLGITPFLNTRTLIRFDDDKGFWC	ÖGLEGGKT	NST-2
172	WGAIRIGVYHKHMKRWLDHFEPIENIHVDGEKLISNPADEISATEKFLGKRVAK	KEKEGVDPPIKKEPCIK	ce3-OST
328	WSAIRIGIYALHLENWLOEPLSÖLLEVSGERLIVDPAGEMARVÖDFTGLKRVVT	KKHEYFNKTKGFPCIKKPEDSSAP	3-OST-4
283	WSAIÖIGIYAKHLEHWRHFEPIRÖMLEVSGERLISDPAGELGRVÖDFTGLKRLIT	DKHEYFNKTKGFPCIKKAEGSSRP	3-OST-3A
245	WNAIRIGMYVLHLESWLÖYFPLAÖIHVSGERLITDPAGEMGRVÖDFTGLKRFIT	DKHEYFNKTKGFPCIKKTESLLP	3-OST-2
188	YKALNRSILYHVHMÖNWLREFPLRHIIHIVDGRILIRDPFEPIÖKVERFLKLSPÖIN	ASNFYFNKTKGFYCLR	3-OST-1
827	KCLGSKRGKYPMDLDSRAFLKDYRDNHIELSKLYKMÖTLPTWLREDLÖNTR		NST-1
826	RCLGRSKGRYPMDTESRLFTLDFERNHNLKSLRSRGÖPVPSPWLREELÖHSSSLG		NST-2
249	HCLGKTKGRHHPDVEPSVLKTLREFYGPENKKEYÖMINHWEDW		ce3-OST
407	RCLGSKGRTHPRIDPDVILRLRKFYKPEINIMFYÖMTGÖDFÖWEÖEEGDK		3-OST-4
362	HCLGKTKGRTHPEIDREVVRRLREFYRPENLKETYÖMTGHDFGWDG		3-OST-3A
324	RCLGSKGRTHVÖIDPEVIDÖLREYFRPYNIKETYETVGÖDFFRWE		3-OST-2
264	RCLHESKGRAHÖVPKLLINKLHEYFHEPNKKEFFELVGRTEFDWH		3-OST-1

Figure 2



EXPRESS MAIL NO.: EL302647095US

PATENT

Attorney Docket No. MIT-087 (5473/91)

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

APPLICANTS: Rosenberg et al.

SERIAL NO.: Not yet assigned      GROUP NO.: Not yet assigned

FILING DATE: Filed herewith      EXAMINER: Not yet assigned

TITLE: HEPARAN SULFATE D-GLUCOSAMINYL 3-O-SULFOTRANSFERASES, AND USES THEREFOR

**BOX PATENT APPLICATION**

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

**SUBMISSION OF PAPER COPY AND/OR COMPUTER READABLE COPY  
OF SEQUENCE LISTING FOR INVENTION  
CONTAINING NUCLEOTIDE AND/OR AMINO ACID SEQUENCE**

(check and complete this item, if applicable)

1. ☐ This replies to the Patent Office Letter dated

Note: If these papers are filed before the office letter issues adequate identification of the original papers should be made, e.g., in addition to the name of the inventor and title of invention, the filing date based on the "Express Mail" procedure, the serial number from the return post card or the attorney's docket number added.

- ☐ A copy of the Patent Office Letter is enclosed.

2. Submitted herewith is

(check each item as applicable)

- A. ☐ a paper copy of the Sequence Listing for this application with each sequence assigned a separate identifier.
- B. ☒ a copy, in computer readable form, of the Sequence Listing for this application.

**STATEMENT**

3. I hereby state that:

(complete applicable items A, B and/or C)

- A. ☒ the content of the paper and computer readable copies submitted herewith are the same.
- B. ☐ the content of the computer readable copy submitted herewith is the same as the Sequence Listing appearing on pages \_\_ to \_\_ of the original specification as filed.
- C. ☒ this submission includes no new matter.

Respectfully submitted,

Date: April 24, 2000  
Reg. No. 44,559

Tel. No.: (617) 248-7103  
Fax No.: (617) 248-7100



Maureen A. Bresnahan  
Attorney for Applicants  
Testa, Hurwitz, & Thibeault, LLP  
High Street Tower  
125 High Street  
Boston, Massachusetts 02110

## SEQUENCE LISTING

<110> ROSENBERG, Robert D  
 SHWORAK, Nicholas W  
 LIU, Jian  
 FRITZE, Linda M. S.  
 SCHWARTZ, John J  
 ZHANG, Lijuan

<120> HEPARAN SULFATE D-GLUCOSAMINYL 3-O-SULFOTRANSFERASES,  
 AND USES THEREFOR

<130> MIT-087

<140>

<141>

<150> WO PCT/US98/22597

<151> 1998-10-23

<150> USSN 60/065,437

<151> 1997-10-31

<150> USSN 60/062,762

<151> 1997-10-24

<160> 16

<170> PatentIn Ver. 2.0

<210> 1

<211> 1685

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (323)..(1255)

<223> mouse 3-OST-1

<400> 1

```

tgcattgcaa tgtgaagtgt tcctgaataa acctgcttga agaaggacaa cgtgggtgttg 60
cgtcttttcc tctgggtcggg gtggaataga cacctccctt ttttaacttg ggtgacctca 120
tgaacataaa agaacttaaa ggtagcaagc catggactta aagtaggctg accttgaact 180
cagagatctt cttggcaatg tctctggaga ttaaagtaat tggcaactgg agatactcat 240
gttccagtaa tcaagaggga gccttgctgc tacttcatga tccaggcgcg tgtggcccag 300
tgaagtccct gagctgtaca gc atg acc ttg ctg ctc ctg ggt gcg gtg ctg 352
Met Thr Leu Leu Leu Leu Gly Ala Val Leu
          1              5              10

ctg gtg gcc cag ccc cag ctt gtg cat tcc cac ccg gct gct cct ggc 400
Leu Val Ala Gln Pro Gln Leu Val His Ser His Pro Ala Ala Pro Gly

```

004240"23243360

				15				20				25							
ccg	ggg	ctc	aaa	cag	cag	gag	ctt	ctg	agg	aag	gtg	att	att	ctc	cca	448			
Pro	Gly	Leu	Lys	Gln	Gln	Glu	Leu	Leu	Arg	Lys	Val	Ile	Ile	Leu	Pro				
30				35				40											
gag	gac	acc	gga	gaa	ggc	aca	gca	tcc	aat	ggg	tcc	aca	cag	cag	ctg	496			
Glu	Asp	Thr	Gly	Glu	Gly	Thr	Ala	Ser	Asn	Gly	Ser	Thr	Gln	Gln	Leu				
45				50				55											
cca	cag	acc	atc	atc	att	ggg	gtg	cgc	aag	ggg	ggg	acc	cga	gcc	ctg	544			
Pro	Gln	Thr	Ile	Ile	Ile	Gly	Val	Arg	Lys	Gly	Gly	Thr	Arg	Ala	Leu				
60				65				70											
cta	gag	atg	ctc	agc	ctg	cat	cct	gat	gtt	gct	gca	gct	gaa	aac	gag	592			
Leu	Glu	Met	Leu	Ser	Leu	His	Pro	Asp	Val	Ala	Ala	Ala	Glu	Asn	Glu				
75				80				85				90							
gtc	cat	ttc	ttt	gac	tgg	gag	gag	cat	tac	agc	caa	ggc	ctg	ggc	tgg	640			
Val	His	Phe	Phe	Asp	Trp	Glu	Glu	His	Tyr	Ser	Gln	Gly	Leu	Gly	Trp				
95				100				105											
tac	ctc	acc	cag	atg	ccc	ttc	tcc	tcc	cct	cac	cag	ctc	acc	gtg	gag	688			
Tyr	Leu	Thr	Gln	Met	Pro	Phe	Ser	Ser	Pro	His	Gln	Leu	Thr	Val	Glu				
110				115				120											
aag	aca	ccc	gcc	tat	ttc	act	tcg	ccc	aaa	gtg	cct	gag	aga	atc	cac	736			
Lys	Thr	Pro	Ala	Tyr	Phe	Thr	Ser	Pro	Lys	Val	Pro	Glu	Arg	Ile	His				
125				130				135											
agc	atg	aac	ccc	acc	atc	cgc	ctg	ctg	ctt	atc	ctg	agg	gac	cca	tca	784			
Ser	Met	Asn	Pro	Thr	Ile	Arg	Leu	Leu	Leu	Ile	Leu	Arg	Asp	Pro	Ser				
140				145				150											
gag	cgc	gtg	ctg	tcc	gac	tac	acc	cag	gtg	ttg	tac	aac	cac	ctt	cag	832			
Glu	Arg	Val	Leu	Ser	Asp	Tyr	Thr	Gln	Val	Leu	Tyr	Asn	His	Leu	Gln				
155				160				165				170							
aag	cac	aag	ccc	tat	cca	ccc	att	gag	gac	ctc	cta	atg	cgg	gac	ggg	880			
Lys	His	Lys	Pro	Tyr	Pro	Pro	Ile	Glu	Asp	Leu	Leu	Met	Arg	Asp	Gly				
175				180				185											
cgg	ctg	aac	ctg	gac	tac	aag	gct	ctc	aac	cgc	agc	ctg	tac	cat	gca	928			
Arg	Leu	Asn	Leu	Asp	Tyr	Lys	Ala	Leu	Asn	Arg	Ser	Leu	Tyr	His	Ala				
190				195				200											
cac	atg	ctg	aac	tgg	ctg	cgt	ttt	ttc	ccg	ttg	ggc	cac	atc	cac	att	976			
His	Met	Leu	Asn	Trp	Leu	Arg	Phe	Phe	Pro	Leu	Gly	His	Ile	His	Ile				
205				210				215											
gtg	gat	ggc	gac	cgc	ctc	atc	aga	gac	cct	ttc	cct	gag	atc	cag	aag	1024			
Val	Asp	Gly	Asp	Arg	Leu	Ile	Arg	Asp	Pro	Phe	Pro	Glu	Ile	Gln	Lys				
220				225				230											
gtc	gaa	aga	ttc	ctg	aag	ctt	tct	cca	cag	atc	aac	gcc	tcg	aac	ttc	1072			
Val	Glu	Arg	Phe	Leu	Lys	Leu	Ser	Pro	Gln	Ile									

00440 "00440" 00440

tac ttt aac aaa acc aag ggc ttc tac tgc ctg cgg gac agt ggc aag 1120  
Tyr Phe Asn Lys Thr Lys Gly Phe Tyr Cys Leu Arg Asp Ser Gly Lys  
255 260 265

gac cgc tgc tta cac gag tcc aaa ggc cgg gcg cac ccc cag gtg gat 1168  
Asp Arg Cys Leu His Glu Ser Lys Gly Arg Ala His Pro Gln Val Asp  
270 275 280

ccc aaa cta ctt gat aaa ctg cac gaa tac ttt cat gag cca aat aag 1216  
Pro Lys Leu Leu Asp Lys Leu His Glu Tyr Phe His Glu Pro Asn Lys  
285 290 295

aaa ttt ttc aag ctc gtg ggc aga aca ttc gac tgg cac tgatttgccg 1265  
Lys Phe Phe Lys Leu Val Gly Arg Thr Phe Asp Trp His  
300 305 310

tctcctaggc tcgggacttt tctgtttgtt aacttctggt gtacatctga aggggggagg 1325

aaaataatth taaaaaggca ttttaagctat aattttatttg taaaaccac aaatgacttc 1385

tgtacagtat tagattcaca gttgccatat atagtagtta tatttttcta cttgttaaat 1445

ggagggcggtt ttgtattgtt tttcatgggtt gttaacattg tgtatatgtc tctataatat 1505

gaaggaactt aactattgca ctgaaaaaat aagagatttt ttttttctgg agacctcttt 1565

ttttgttggtt gttgttttaa atataattaa cctgcctcca atccaaaata gctctttgtt 1625

ttcacctcct tgtcaaactc ataactctttt totgcttaaa aaatttattg gtattatgga 1685

<210> 2

<211> 311

<212> PRT

<213> Mus musculus

<400> 2

Met Thr Leu Leu Leu Leu Gly Ala Val Leu Leu Val Ala Gln Pro Gln  
1 5 10 15

Leu Val His Ser His Pro Ala Ala Pro Gly Pro Gly Leu Lys Gln Gln  
20 25 30

Glu Leu Leu Arg Lys Val Ile Ile Leu Pro Glu Asp Thr Gly Glu Gly  
35 40 45

Thr Ala Ser Asn Gly Ser Thr Gln Gln Leu Pro Gln Thr Ile Ile Ile  
50 55 60

Gly Val Arg Lys Gly Gly Thr Arg Ala Leu Leu Glu Met Leu Ser Leu  
65 70 75 80

His Pro Asp Val Ala Ala Ala Glu Asn Glu Val His Phe Phe Asp Trp  
85 90 95

Glu Glu His Tyr Ser Gln Gly Leu Gly Trp Tyr Leu Thr Gln Met Pro



004240"2924350

100					105					110					
Phe	Ser	Ser	Pro	His	Gln	Leu	Thr	Val	Glu	Lys	Thr	Pro	Ala	Tyr	Phe
		115					120					125			
Thr	Ser	Pro	Lys	Val	Pro	Glu	Arg	Ile	His	Ser	Met	Asn	Pro	Thr	Ile
	130					135					140				
Arg	Leu	Leu	Leu	Ile	Leu	Arg	Asp	Pro	Ser	Glu	Arg	Val	Leu	Ser	Asp
145					150					155					160
Tyr	Thr	Gln	Val	Leu	Tyr	Asn	His	Leu	Gln	Lys	His	Lys	Pro	Tyr	Pro
			165						170					175	
Pro	Ile	Glu	Asp	Leu	Leu	Met	Arg	Asp	Gly	Arg	Leu	Asn	Leu	Asp	Tyr
			180					185					190		
Lys	Ala	Leu	Asn	Arg	Ser	Leu	Tyr	His	Ala	His	Met	Leu	Asn	Trp	Leu
		195					200					205			
Arg	Phe	Phe	Pro	Leu	Gly	His	Ile	His	Ile	Val	Asp	Gly	Asp	Arg	Leu
	210					215						220			
Ile	Arg	Asp	Pro	Phe	Pro	Glu	Ile	Gln	Lys	Val	Glu	Arg	Phe	Leu	Lys
225					230					235					240
Leu	Ser	Pro	Gln	Ile	Asn	Ala	Ser	Asn	Phe	Tyr	Phe	Asn	Lys	Thr	Lys
				245					250					255	
Gly	Phe	Tyr	Cys	Leu	Arg	Asp	Ser	Gly	Lys	Asp	Arg	Cys	Leu	His	Glu
			260					265					270		
Ser	Lys	Gly	Arg	Ala	His	Pro	Gln	Val	Asp	Pro	Lys	Leu	Leu	Asp	Lys
		275					280					285			
Leu	His	Glu	Tyr	Phe	His	Glu	Pro	Asn	Lys	Lys	Phe	Phe	Lys	Leu	Val
	290					295									
Gly	Arg	Thr	Phe	Asp	Trp	His									
305					310										

<210> 3  
 <211> 1305  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (119)..(1039)  
 <223> human 3-OST-1

<400> 3  
 cgcggtctcag taattgaagg cctgaaacgc ccatgtgcc a ctgactagga ggcttcctg 60  
 ctgcggcact tcatgaccca gcggcgcgcg gcccagtga gccaccgtgg tgtccagc 118

atg gcc gcg ctg ctc ctg ggc gcg gtg ctg ctg gtg gcc cag ccc cag	166
Met Ala Ala Leu Leu Leu Gly Ala Val Leu Leu Val Ala Gln Pro Gln	
1 5 10 15	
cta gtg cct tcc cgc ccc gcc gag cta ggc cag cag gag ctt ctg cgg	214
Leu Val Pro Ser Arg Pro Ala Glu Leu Gly Gln Gln Glu Leu Leu Arg	
20 25 30	
aaa gcg ggg acc ctc cag gat gac gtc cgc gat ggc gtg gcc cca aac	262
Lys Ala Gly Thr Leu Gln Asp Asp Val Arg Asp Gly Val Ala Pro Asn	
35 40 45	
ggc tct gcc cag cag ttg ccg cag acc atc atc atc ggc gtg cgc aag	310
Gly Ser Ala Gln Gln Leu Pro Gln Thr Ile Ile Ile Gly Val Arg Lys	
50 55 60	
ggc ggc acg cgc gca ctg ctg gag atg ctc agc ctg cac ccc gac gtg	358
Gly Gly Thr Arg Ala Leu Leu Glu Met Leu Ser Leu His Pro Asp Val	
65 70 75 80	
gcg gcc gcg gag aac gag gtc cac ttc ttc gac tgg gag gag cat tac	406
Ala Ala Ala Glu Asn Glu Val His Phe Phe Asp Trp Glu Glu His Tyr	
85 90 95	
agc cac ggc ttg ggc tgg tac ctc agc cag atg ccc ttc tcc tgg cca	454
Ser His Gly Leu Gly Trp Tyr Leu Ser Gln Met Pro Phe Ser Trp Pro	
100 105 110	
cac cag ctc aca gtg gag aag acc ccc gcg tat ttc acg tcg ccc aaa	502
His Gln Leu Thr Val Glu Lys Thr Pro Ala Tyr Phe Thr Ser Pro Lys	
115 120 125	
gtg cct gag cga gtc tac agc atg aac ccg tcc atc cgg ctg ctg ctc	550
Val Pro Glu Arg Val Tyr Ser Met Asn Pro Ser Ile Arg Leu Leu Leu	
130 135 140	
atc ctg cga gac ccg tcg gag cgc gtg cta tct gac tac acc caa gtg	598
Ile Leu Arg Asp Pro Ser Glu Arg Val Leu Ser Asp Tyr Thr Gln Val	
145 150 155 160	
ttc tac aac cac atg cag aag cac aag ccc tac ccg tcc atc gag gag	646
Phe Tyr Asn His Met Gln Lys His Lys Pro Tyr Pro Ser Ile Glu Glu	
165 170 175	
ttc ctg gtg cgc gat ggc agg ctc aat gtg gac tac aag gcc ctc aac	694
Phe Leu Val Arg Asp Gly Arg Leu Asn Val Asp Tyr Lys Ala Leu Asn	
180 185 190	
cgc agc ctc tac cac gtg cac atg cag aac tgg ctg cgc ttt ttc ccg	742
Arg Ser Leu Tyr His Val His Met Gln Asn Trp Leu Arg Phe Phe Pro	
195 200 205	
ctg cgc cac atc cac att gtg gac ggc gac cgc ctc atc agg gac ccc	790
Leu Arg His Ile His Ile Val Asp Gly Asp Arg Leu Ile Arg Asp Pro	
210 215 220	
ttc cct gag atc caa aag gtc gag agg ttc cta aag ctg tcg ccg cag	838

```

Phe Pro Glu Ile Gln Lys Val Glu Arg Phe Leu Lys Leu Ser Pro Gln
225                230                235                240

atc aat gct tcg aac ttc tac ttt aac aaa acc aag ggc ttt tac tgc   886
Ile Asn Ala Ser Asn Phe Tyr Phe Asn Lys Thr Lys Gly Phe Tyr Cys
                245                250                255

ctg cgg gac agc ggc cgg gac cgc tgc tta cat gag tcc aaa ggc cgg   934
Leu Arg Asp Ser Gly Arg Asp Arg Cys Leu His Glu Ser Lys Gly Arg
                260                265                270

gcg cac ccc caa gtc gat ccc aaa cta ctc aat aaa ctg cac gaa tat   982
Ala His Pro Gln Val Asp Pro Lys Leu Leu Asn Lys Leu His Glu Tyr
                275                280                285

ttt cat gag cca aat aag aag ttc ttc gag ctt gtt ggc aga aca ttt   1030
Phe His Glu Pro Asn Lys Lys Phe Phe Glu Leu Val Gly Arg Thr Phe
                290                295                300

gac tgg cac tgatttgcaa taagctaagc tcagaaactt tcctactgta           1079
Asp Trp His
305

agttctggtg tacatctgag gggaaaaaga attttaaaaa agcattttaag gtataattta 1139

tttgtaaaat ccataaagta cttctgtaca gtattagatt cacaattgcc atatatacta 1199

gttatatttt tctacttggt aaatggaggg cattttgtat tgtttttcat ggttggttaac 1259

attgtgtaat atgtctctat atgaaggaac taaactattt cactga                1305

<210> 4
<211> 307
<212> PRT
<213> Homo sapiens

<400> 4
Met Ala Ala Leu Leu Leu Gly Ala Val Leu Leu Val Ala Gln Pro Gln
  1                5                10                15

Leu Val Pro Ser Arg Pro Ala Glu Leu Gly Gln Gln Glu Leu Leu Arg
                20                25                30

Lys Ala Gly Thr Leu Gln Asp Asp Val Arg Asp Gly Val Ala Pro Asn
  35                40                45

Gly Ser Ala Gln Gln Leu Pro Gln Thr Ile Ile Ile Gly Val Arg Lys
  50                55                60

Gly Gly Thr Arg Ala Leu Leu Glu Met Leu Ser Leu His Pro Asp Val
  65                70                75                80

Ala Ala Ala Glu Asn Glu Val His Phe Phe Asp Trp Glu Glu His Tyr
                85                90                95

Ser His Gly Leu Gly Trp Tyr Leu Ser Gln Met Pro Phe Ser Trp Pro

```

100							105					110			
His	Gln	Leu	Thr	Val	Glu	Lys	Thr	Pro	Ala	Tyr	Phe	Thr	Ser	Pro	Lys
		115				120				125					
Val	Pro	Glu	Arg	Val	Tyr	Ser	Met	Asn	Pro	Ser	Ile	Arg	Leu	Leu	Leu
130						135				140					
Ile	Leu	Arg	Asp	Pro	Ser	Glu	Arg	Val	Leu	Ser	Asp	Tyr	Thr	Gln	Val
145				150						155				160	
Phe	Tyr	Asn	His	Met	Gln	Lys	His	Lys	Pro	Tyr	Pro	Ser	Ile	Glu	Glu
				165				170						175	
Phe	Leu	Val	Arg	Asp	Gly	Arg	Leu	Asn	Val	Asp	Tyr	Lys	Ala	Leu	Asn
		180						185				190			
Arg	Ser	Leu	Tyr	His	Val	His	Met	Gln	Asn	Trp	Leu	Arg	Phe	Phe	Pro
		195				200						205			
Leu	Arg	His	Ile	His	Ile	Val	Asp	Gly	Asp	Arg	Leu	Ile	Arg	Asp	Pro
210						215				220					
Phe	Pro	Glu	Ile	Gln	Lys	Val	Glu	Arg	Phe	Leu	Lys	Leu	Ser	Pro	Gln
225				230						235				240	
Ile	Asn	Ala	Ser	Asn	Phe	Tyr	Phe	Asn	Lys	Thr	Lys	Gly	Phe	Tyr	Cys
				245				250						255	
Leu	Arg	Asp	Ser	Gly	Arg	Asp	Arg	Cys	Leu	His	Glu	Ser	Lys	Gly	Arg
		260						265				270			
Ala	His	Pro	Gln	Val	Asp	Pro	Lys	Leu	Leu	Asn	Lys	Leu	His	Glu	Tyr
		275				280						285			
Phe	His	Glu	Pro	Asn	Lys	Lys	Phe	Phe	Glu	Leu	Val	Gly	Arg	Thr	Phe
290						295				300					
Asp	Trp	His													
305															

```
<210> 5
<211> 1951
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> CDS
<222> (73)..(1173)
<223> human 3-OST-2
```

<400> 5  
cgcagggccca cagcagctca gccgcgggtg cccctctcga aaccatgacc cccqgcgcgq 60

gcccatggag cc atg gcc tat agg gtc ctg ggc cgc gcg ggg cca cct cag 111  
Met Ala Tyr Arg Val Leu Gly Arg Ala Gly Pro Pro Gln

			1				5				10							
ccg	cgg	agg	gcg	cgc	agg	ctg	ctc	ttc	gcc	ttc	acg	ctc	tgc	ctc	tcc	159		
Pro	Arg	Arg	Ala	Arg	Arg	Leu	Leu	Phe	Ala	Phe	Thr	Leu	Ser	Leu	Ser			
			15			20			25									
tgc	act	tac	ctg	tgt	tac	agc	ttc	ctg	tgc	tgc	tgc	gac	gac	ctg	ggc	207		
Cys	Thr	Tyr	Leu	Cys	Tyr	Ser	Phe	Leu	Cys	Cys	Cys	Asp	Asp	Leu	Gly			
			30			35			40			45						
cgg	agc	cgc	ctc	ctc	ggc	gcg	cct	cgc	tgc	ctc	cgc	ggc	ccc	agc	gcg	255		
Arg	Ser	Arg	Leu	Leu	Gly	Ala	Pro	Arg	Cys	Leu	Arg	Gly	Pro	Ser	Ala			
			50			55			60									
ggc	ggc	cag	aaa	ctt	ctc	cag	aag	tcc	cgc	ccc	tgt	gat	ccc	tcc	ggg	303		
Gly	Gly	Gln	Lys	Leu	Leu	Gln	Lys	Ser	Arg	Pro	Cys	Asp	Pro	Ser	Gly			
			65			70			75									
ccg	acg	ccc	agc	gag	ccc	agc	gct	ccc	agc	gcg	ccc	gcc	gcc	gcc	gtg	351		
Pro	Thr	Pro	Ser	Glu	Pro	Ser	Ala	Pro	Ser	Ala	Pro	Ala	Ala	Ala	Val			
			80			85			90									
ccc	gcc	cct	cgc	ctc	tcc	ggc	tcc	aac	cac	tcc	ggc	tca	ccc	aag	ctg	399		
Pro	Ala	Pro	Arg	Leu	Ser	Gly	Ser	Asn	His	Ser	Gly	Ser	Pro	Lys	Leu			
			95			100			105									
ggc	acc	aag	cgg	ttg	ccc	caa	gcc	ctc	att	gtg	ggc	gtg	aag	aag	ggg	447		
Gly	Thr	Lys	Arg	Leu	Pro	Gln	Ala	Leu	Ile	Val	Gly	Val	Lys	Lys	Gly			
			110			115			120			125						
ggc	acc	cgg	gcc	gtg	ctg	gag	ttt	atc	cga	gta	cac	ccg	gac	gtg	cgg	495		
Gly	Thr	Arg	Ala	Val	Leu	Glu	Phe	Ile	Arg	Val	His	Pro	Asp	Val	Arg			
			130			135			140									
gcc	ttg	ggc	acg	gaa	ccc	cac	ttc	ttt	gac	agg	aac	tac	ggc	cgc	ggg	543		
Ala	Leu	Gly	Thr	Glu	Pro	His	Phe	Phe	Asp	Arg	Asn	Tyr	Gly	Arg	Gly			
			145			150			155									
ctg	gat	tgg	tac	agg	agc	ctg	atg	ccc	agg	acc	ctc	gag	agc	cag	atc	591		
Leu	Asp	Trp	Tyr	Arg	Ser	Leu	Met	Pro	Arg	Thr	Leu	Glu	Ser	Gln	Ile			
			160			165			170									
acg	ctg	gag	aag	acg	ccc	agc	tac	ttt	gtc	act	caa	gag	gct	cct	cga	639		
Thr	Leu	Glu	Lys	Thr	Pro	Ser	Tyr	Phe	Val	Thr	Gln	Glu	Ala	Pro	Arg			
			175			180			185									
cgc	atc	ttc	aac	atg	tcc	cga	gac	acc	aag	ctg	atc	gtg	gtt	gtg	cgg	687		
Arg	Ile	Phe	Asn	Met	Ser	Arg	Asp	Thr	Lys	Leu	Ile	Val	Val	Val	Arg			
			190			195			200			205						
aac	cct	gtg	acc	cgt	gcc	atc	tct	gat	tac	acg	cag	aca	ctc	tcc	aag	735		
Asn	Pro	Val	Thr	Arg	Ala	Ile	Ser	Asp	Tyr	Thr	Gln	Thr	Leu	Ser	Lys			
			210			215			220									
aag	ccc	gac	atc	ccg	acc	ttt	gag	ggc	ctc	tcc	ttc	cgc	aac	cgc	acc	783		
Lys	Pro	Asp	Ile	Pro	Thr	Phe	Glu	Gly	Leu	Ser	Phe	Arg	Asn	Arg	Thr			
			225			230			235									

ctg ggc ctg gtg gac gtg tcg tgg aac gcc atc cgc atc ggc atg tac	831
Leu Gly Leu Val Asp Val Ser Trp Asn Ala Ile Arg Ile Gly Met Tyr	
240 245 250	
gtg ctg cac ctg gag agc tgg ctg cag tac ttc ccg cta gct cag att	879
Val Leu His Leu Glu Ser Trp Leu Gln Tyr Phe Pro Leu Ala Gln Ile	
255 260 265	
cac ttc gtc agt ggc gag cga ctc atc act gac ccg gcc ggc gag atg	927
His Phe Val Ser Gly Glu Arg Leu Ile Thr Asp Pro Ala Gly Glu Met	
270 275 280 285	
ggg cga gtc cag gac ttc ctg ggc att aag aga ttc atc acg gac aag	975
Gly Arg Val Gln Asp Phe Leu Gly Ile Lys Arg Phe Ile Thr Asp Lys	
290 295 300	
cac ttc tat ttc aac aag acc aaa gga ttc cct tgc ttg aaa aaa aca	1023
His Phe Tyr Phe Asn Lys Thr Lys Gly Phe Pro Cys Leu Lys Lys Thr	
305 310 315	
gaa tcg agc ctc ctg cct cga tgc ttg ggc aaa tca aaa ggg aga act	1071
Glu Ser Ser Leu Leu Pro Arg Cys Leu Gly Lys Ser Lys Gly Arg Thr	
320 325 330	
cat gta cag att gat cct gaa gtg ata gac cag ctc cga gaa ttt tat	1119
His Val Gln Ile Asp Pro Glu Val Ile Asp Gln Leu Arg Glu Phe Tyr	
335 340 345	
aga ccg tat aat atc aaa ttt tat gaa acc gtt ggg cag gac ttc agg	1167
Arg Pro Tyr Asn Ile Lys Phe Tyr Glu Thr Val Gly Gln Asp Phe Arg	
350 355 360 365	
tgg gaa taagcccacg aaaggaaagg gctctcaagg gctcttctgc tcattctctc	1223
Trp Glu	
cgtagagattt gctccagac cctcttatct cctccaaca aaccctggct ccagccccct	1283
ttcccaactt gagttgcac atcttggaac caggaagccc agctaaagcc aagagaccag	1343
agagtctctg ccactagttt tcattcagtt gttcaagcaa agttgatctg ctctggcac	1403
gtccagtaaa ttccagaatc attctccttt ctgccataa agggccttgg agaattgctt	1463
taagaagagt gaatgttcca atgatgatag atattataag cgacgatggg tctgttgcta	1523
tgaacacagc agtcgggtccc tgtcattgtc caccaggag tggccttggt aattccaagt	1583
ggcatgtatc ttccctctga gcttcatttc ttcaagatgc tctgggtggg gggatgggag	1643
accatcctca gccctcctca gaccttatca attcattgag agattgcaaa gctgaaagca	1703
cctccggcca ctctggggag acagaccctt tggatgatgaa ataaaccagt gacttcagag	1763
cctatgggtct caactgtgct tgaaaaacac tgtctctgaa aacaactttg tgattctccc	1823
tgctccctgt ggacaaaagc acataattct gctgttacgg gtactttgct catacgagct	1883

ttcatgttca gcatgcaatg gaatcatgct tgtccatgtg aaataaatat ggctctctcg 1943

tgtcctta 1951

<210> 6

<211> 367

<212> PRT

<213> Homo sapiens

<400> 6

Met Ala Tyr Arg Val Leu Gly Arg Ala Gly Pro Pro Gln Pro Arg Arg  
1 5 10 15

Ala Arg Arg Leu Leu Phe Ala Phe Thr Leu Ser Leu Ser Cys Thr Tyr  
20 25 30

Leu Cys Tyr Ser Phe Leu Cys Cys Cys Asp Asp Leu Gly Arg Ser Arg  
35 40 45

Leu Leu Gly Ala Pro Arg Cys Leu Arg Gly Pro Ser Ala Gly Gly Gln  
50 55 60

Lys Leu Leu Gln Lys Ser Arg Pro Cys Asp Pro Ser Gly Pro Thr Pro  
65 70 75 80

Ser Glu Pro Ser Ala Pro Ser Ala Pro Ala Ala Val Pro Ala Pro  
85 90 95

Arg Leu Ser Gly Ser Asn His Ser Gly Ser Pro Lys Leu Gly Thr Lys  
100 105 110

Arg Leu Pro Gln Ala Leu Ile Val Gly Val Lys Lys Gly Gly Thr Arg  
115 120 125

Ala Val Leu Glu Phe Ile Arg Val His Pro Asp Val Arg Ala Leu Gly  
130 135 140

Thr Glu Pro His Phe Phe Asp Arg Asn Tyr Gly Arg Gly Leu Asp Trp  
145 150 155 160

Tyr Arg Ser Leu Met Pro Arg Thr Leu Glu Ser Gln Ile Thr Leu Glu  
165 170 175

Lys Thr Pro Ser Tyr Phe Val Thr Gln Glu Ala Pro Arg Arg Ile Phe  
180 185 190

Asn Met Ser Arg Asp Thr Lys Leu Ile Val Val Val Arg Asn Pro Val  
195 200 205

Thr Arg Ala Ile Ser Asp Tyr Thr Gln Thr Leu Ser Lys Lys Pro Asp  
210 215 220

Ile Pro Thr Phe Glu Gly Leu Ser Phe Arg Asn Arg Thr Leu Gly Leu  
225 230 235 240

bioRxiv preprint doi: <https://doi.org/10.1101/2024.08.28.610560>; this version posted September 3, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Val Asp Val Ser Trp Asn Ala Ile Arg Ile Gly Met Tyr Val Leu His  
245 250 255

Leu Glu Ser Trp Leu Gln Tyr Phe Pro Leu Ala Gln Ile His Phe Val  
260 265 270

Ser Gly Glu Arg Leu Ile Thr Asp Pro Ala Gly Glu Met Gly Arg Val  
275 280 285

Gln Asp Phe Leu Gly Ile Lys Arg Phe Ile Thr Asp Lys His Phe Tyr  
290 295 300

Phe Asn Lys Thr Lys Gly Phe Pro Cys Leu Lys Lys Thr Glu Ser Ser  
305 310 315 320

Leu Leu Pro Arg Cys Leu Gly Lys Ser Lys Gly Arg Thr His Val Gln  
325 330 335

Ile Asp Pro Glu Val Ile Asp Gln Leu Arg Glu Phe Tyr Arg Pro Tyr  
340 345 350

Asn Ile Lys Phe Tyr Glu Thr Val Gly Gln Asp Phe Arg Trp Glu  
355 360 365

<210> 7  
<211> 2314  
<212> DNA  
<213> Homo sapiens

<220>  
<221> CDS  
<222> (799) .. (2016)  
<223> human 3-OST-3A

<400> 7  
cagcggcggc ccaggaggca gccggtgagc gcctgcgagc agagtggcgg gggccgctga 60  
caggtcccg cgcagcccagc ccagcccagc cagcgggctc acaggtgggg tccaagagca 120  
gtttggagca acccggcgct acggagaggg gtggacggct ctgcacgggc ctctgtctc 180  
ccgctcgggc agaggggactc ggggggacct cgctccttgg ccgagagAAC ctgaactcgg 240  
gcggagagaa cgcgcccagg cgggcaaggg gaccagagaa agccggggct ggaagtcact 300  
gtcgctcgcc actgtctgga ggcacaggag cgcagaggcc cggcagccgc gcgtgccctc 360  
ccggggaccg agccagtgat gcaggatcgc tgagcggaga tccgcgccga gaagtctctc 420  
ggggccgggg ctgagacgca cgccttcgac accgctgcc agaccccgat tccggcgact 480  
cttcggggga accgaggggc caaggctgcc ccaagctcag gacttgggcg agtctaagac 540  
gatggtttct taagcacgga ccgcggttcc ccttcccgcc cctcgactg gaggcaggga 600  
tcctgcgcgg gggccccggg attccgtttc ccgcgggagc ccggccgct gcctcccggg 660



acagttcgca	cggccacagg	ggcgcacggc	gatgtggcct	ccgtccagcg	cgctggccccg	720
ccgggggggat	gctctggcac	ctgtcggggg	ccaggcctag	catggccggc	gcgttgccccg	780
acgtcgccctc	cggtctagg	atg gcc cct ccg ggc ccg gcc agt gcc ctc tcc	831			
		Met Ala Pro Pro Gly Pro Ala Ser Ala Leu Ser				
		1 5 10				
acc tcg gcc gag ccg ctg tcc cgc agc atc ttc ccg aag ttc ttg ctg	879					
Thr Ser Ala Glu Pro Leu Ser Arg Ser Ile Phe Arg Lys Phe Leu Leu						
	15 20 25					
atg ctc tgc tcc ctg ctc acg tcc ctt tac gtc ttc tac tgc ctg gcc	927					
Met Leu Cys Ser Leu Leu Thr Ser Leu Tyr Val Phe Tyr Cys Leu Ala						
	30 35 40					
gag cgc tgc cag acc ctg tcc ggc ccc gtc gtg ggg ctg tcc ggc ggc	975					
Glu Arg Cys Gln Thr Leu Ser Gly Pro Val Val Gly Leu Ser Gly Gly						
	45 50 55					
ggc gag gag gcg ggg gcc cct ggt ggc ggc gtc ctg gcc gga ggc ccg	1023					
Gly Glu Glu Ala Gly Ala Pro Gly Gly Gly Val Leu Ala Gly Gly Pro						
	60 65 70 75					
agg gag ctg gcg gtg tgg ccg gcg gcg gca cag aga aag cgc ctc ctg	1071					
Arg Glu Leu Ala Val Trp Pro Ala Ala Ala Gln Arg Lys Arg Leu Leu						
	80 85 90					
caa ctg ccg cag tgg ccg agg cgc ccg ccg ccc gcg ccc cgc gac gac	1119					
Gln Leu Pro Gln Trp Arg Arg Arg Arg Pro Pro Ala Pro Arg Asp Asp						
	95 100 105					
ggc gag gag gcg gcc tgg gaa gaa gag tcc cct ggc ctg tca ggg ggt	1167					
Gly Glu Glu Ala Ala Trp Glu Glu Glu Ser Pro Gly Leu Ser Gly Gly						
	110 115 120					
ccg ggc ggc tcc ggg gcc gga agc acc gtg gcc gag gcc ccg ccg ggg	1215					
Pro Gly Gly Ser Gly Ala Gly Ser Thr Val Ala Glu Ala Pro Pro Gly						
	125 130 135					
acc ctg gcg ctg ctc ctg gac gaa ggc agc aag cag ctg ccg cag gcc	1263					
Thr Leu Ala Leu Leu Leu Asp Glu Gly Ser Lys Gln Leu Pro Gln Ala						
	140 145 150 155					
atc atc atc gga gtg aag aag ggc ggc acg ccg gcg ctg ctg gag ttc	1311					
Ile Ile Ile Gly Val Lys Lys Gly Gly Thr Arg Ala Leu Leu Glu Phe						
	160 165 170					
ctg cgc gtg cac ccc gac gtg cgc gcc gtg ggc gcc gag ccc cac ttc	1359					
Leu Arg Val His Pro Asp Val Arg Ala Val Gly Ala Glu Pro His Phe						
	175 180 185					
ttc gac cgc agc tac gac aag ggc ctc gcc tgg tac ccg gac ctg atg	1407					
Phe Asp Arg Ser Tyr Asp Lys Gly Leu Ala Trp Tyr Arg Asp Leu Met						
	190 195 200					

ccc aga acc ctg gac ggg cag atc acc atg gag aag acg ccc agt tac 1455  
Pro Arg Thr Leu Asp Gly Gln Ile Thr Met Glu Lys Thr Pro Ser Tyr  
205 210 215

ttc gtc acg cgg gag gcc ccc gcg cgc atc tcg gcc atg tcc aag gac 1503  
Phe Val Thr Arg Glu Ala Pro Ala Arg Ile Ser Ala Met Ser Lys Asp  
220 225 230 235

acc aag ctc atc gtg gtg gtg cgg gac ccg gtg acc agg gcc atc tcg 1551  
Thr Lys Leu Ile Val Val Val Arg Asp Pro Val Thr Arg Ala Ile Ser  
240 245 250

gac tac acg cag acg ctg tcc aag cgg ccc gac atc ccc acc ttc gag 1599  
Asp Tyr Thr Gln Thr Leu Ser Lys Arg Pro Asp Ile Pro Thr Phe Glu  
255 260 265

agc ttg acg ttc aaa aac agg aca gcg ggc ctc atc gac acg tcg tgg 1647  
Ser Leu Thr Phe Lys Asn Arg Thr Ala Gly Leu Ile Asp Thr Ser Trp  
270 275 280

agc gcc atc cag atc ggc atc tac gcc aag cac ctg gag cac tgg ctg 1695  
Ser Ala Ile Gln Ile Gly Ile Tyr Ala Lys His Leu Glu His Trp Leu  
285 290 295

cgc cac ttc ccc atc cgc cag atg ctc ttc gtg agc ggc gag cgg ctc 1743  
Arg His Phe Pro Ile Arg Gln Met Leu Phe Val Ser Gly Glu Arg Leu  
300 305 310 315

atc agc gac ccg gcc ggg gag ctg ggc cgc gtg caa gac ttc ctg ggc 1791  
Ile Ser Asp Pro Ala Gly Glu Leu Gly Arg Val Gln Asp Phe Leu Gly  
320 325 330

ctc aag agg atc atc acg gac aag cac ttc tac ttc aac aag acc aag 1839  
Leu Lys Arg Ile Ile Thr Asp Lys His Phe Tyr Phe Asn Lys Thr Lys  
335 340 345

ggc ttc ccc tgc ctg aag aag gcg gag ggc agc agc cgg ccc cat tgc 1887  
Gly Phe Pro Cys Leu Lys Lys Ala Glu Gly Ser Ser Arg Pro His Cys  
350 355 360

ctg ggc aag acc aag ggc agg acc cat cct gag atc gac cgc gag gtg 1935  
Leu Gly Lys Thr Lys Gly Arg Thr His Pro Glu Ile Asp Arg Glu Val  
365 370 375

gtg cgc agg ctg cgc gag ttc tac cgg cct ttc aac ctc aag ttc tac 1983  
Val Arg Arg Leu Arg Glu Phe Tyr Arg Pro Phe Asn Leu Lys Phe Tyr  
380 385 390 395

cag atg acc ggg cac gac ttt ggc tgg gat gga taaccatata atttaaaaag 2036  
Gln Met Thr Gly His Asp Phe Gly Trp Asp Gly  
400 405

aaaaaaaaaa tcaaaatata atatattttt ttaccaatcg gtagagaaga gacagtttaa 2096

tatttggtgct gaaaatatgt ttcagtattt ttttcaatga atgttaagag attgtttctca 2156

ctcccgcccc atcttaatgt ataaccaaca ccaaacacgt ggatcaacag aaaaggaaaa 2216



Val Val Arg Asp Pro Val Thr Arg Ala Ile Ser Asp Tyr Thr Gln Thr  
245 250 255

Leu Ser Lys Arg Pro Asp Ile Pro Thr Phe Glu Ser Leu Thr Phe Lys  
260 265 270

Asn Arg Thr Ala Gly Leu Ile Asp Thr Ser Trp Ser Ala Ile Gln Ile  
275 280 285

Gly Ile Tyr Ala Lys His Leu Glu His Trp Leu Arg His Phe Pro Ile  
290 295 300

Arg Gln Met Leu Phe Val Ser Gly Glu Arg Leu Ile Ser Asp Pro Ala  
305 310 315 320

Gly Glu Leu Gly Arg Val Gln Asp Phe Leu Gly Leu Lys Arg Ile Ile  
325 330 335

Thr Asp Lys His Phe Tyr Phe Asn Lys Thr Lys Gly Phe Pro Cys Leu  
340 345 350

Lys Lys Ala Glu Gly Ser Ser Arg Pro His Cys Leu Gly Lys Thr Lys  
355 360 365

Gly Arg Thr His Pro Glu Ile Asp Arg Glu Val Val Arg Arg Leu Arg  
370 375 380

Glu Phe Tyr Arg Pro Phe Asn Leu Lys Phe Tyr Gln Met Thr Gly His  
385 390 395 400

Asp Phe Gly Trp Asp Gly  
405

<210> 9  
<211> 2032  
<212> DNA  
<213> Homo sapiens

<220>  
<221> CDS  
<222> (331) .. (1500)  
<223> human 3-OST-3B

<400> 9  
gtggccaggg cgcgagagtg caacgtcctc ctggccccga gcgcgtcgtc gcgccccggg 60  
agcagaccct cgcccagcag ttaccgccgt cccgactttc cgttccagtt gcagctcctg 120  
ccgggcaaca tgtcaagagc cgccgccgct acagctgccg ccgccacctg gggaagagca 180  
gcagcagcag cggcgggccgc gggcacacgg gggcaataaa ccgagccacc cgggcgtcca 240  
gcgtgccggg gaaccctctc tgcgtcact gcccggcggg acccacgcca tgtgctgagc 300  
catgtccctg gccgcgcccg cgggcagcgc atg ggg cag cgc ctg agt ggc ggc 354  
Met Gly Gln Arg Leu Ser Gly Gly

															1																5	
aga	tct	tgc	ctc	gat	gtc	ccc	ggc	cgg	ctc	cta	ccg	cag	ccg	ccg	ccg	402																
Arg	Ser	Cys	Leu	Asp	Val	Pro	Gly	Arg	Leu	Leu	Pro	Gln	Pro	Pro	Pro																	
					10						15						20															
ccc	ccg	ccg	ccg	gtg	agg	agg	aag	ctc	gcg	ctg	ctc	ttc	gcc	atg	ctc	450																
Pro	Pro	Pro	Pro	Val	Arg	Arg	Lys	Leu	Ala	Leu	Leu	Phe	Ala	Met	Leu																	
					25						30						35	40														
tgc	gtc	tgg	ctc	tat	atg	ttc	ctg	tac	tgc	tgc	gcc	ggc	tcc	tgc	gcc	498																
Cys	Val	Trp	Leu	Tyr	Met	Phe	Leu	Tyr	Ser	Cys	Ala	Gly	Ser	Cys	Ala																	
					45						50						55															
gcc	gcg	ccg	ggg	ctg	ctg	ctc	ctg	ggc	tct	ggg	tcc	cgc	gcc	gca	cac	546																
Ala	Ala	Pro	Gly	Leu	Leu	Leu	Leu	Gly	Ser	Gly	Ser	Arg	Ala	Ala	His																	
					60						65						70															
gac	ccg	cca	gcc	ctg	gcc	aca	gct	ccg	gac	ggg	acg	ccc	ccc	agg	ctg	594																
Asp	Pro	Pro	Ala	Leu	Ala	Thr	Ala	Pro	Asp	Gly	Thr	Pro	Pro	Arg	Leu																	
					75						80						85															
ccg	ttc	egg	gcg	ccg	cca	gcc	acc	cca	ctg	gct	tca	ggc	aag	gag	atg	642																
Pro	Phe	Arg	Ala	Pro	Pro	Ala	Thr	Pro	Leu	Ala	Ser	Gly	Lys	Glu	Met																	
					90						95						100															
gcc	gag	ggc	gct	gcg	agc	ccg	gag	gag	cag	agt	ccc	gag	gtg	ccg	gac	690																
Ala	Glu	Gly	Ala	Ala	Ser	Pro	Glu	Glu	Gln	Ser	Pro	Glu	Val	Pro	Asp																	
					105						110						115	120														
tcc	cca	agc	ccc	atc	tcc	agc	ttt	ttc	agt	ggg	tct	ggg	agc	aag	cag	738																
Ser	Pro	Ser	Pro	Ile	Ser	Ser	Phe	Phe	Ser	Gly	Ser	Gly	Ser	Lys	Gln																	
					125						130						135															
ctg	ccg	cag	gcc	atc	atc	atc	ggc	gtg	aag	aag	ggc	ggc	acg	cgg	gcg	786																
Leu	Pro	Gln	Ala	Ile	Ile	Ile	Gly	Val	Lys	Lys	Gly	Gly	Thr	Arg	Ala																	
					140						145						150															
ctg	ctg	gag	ttt	ctg	cgc	gtg	cac	ccc	gac	gtg	cgc	gcc	gtg	ggc	gcc	834																
Leu	Leu	Glu	Phe	Leu	Arg	Val	His	Pro	Asp	Val	Arg	Ala	Val	Gly	Ala																	
					155						160						165															
gag	ccc	cat	ttc	ttc	gat	cgc	agc	tac	gac	aag	ggc	ctc	gct	tgg	tac	882																
Glu	Pro	His	Phe	Phe	Asp	Arg	Ser	Tyr	Asp	Lys	Gly	Leu	Ala	Trp	Tyr																	
					170						175						180															
cgg	gac	ctg	atg	ccc	aga	acc	ctg	gac	ggg	cag	atc	acc	atg	gag	aag	930																
Arg	Asp	Leu	Met	Pro	Arg	Thr	Leu	Asp	Gly	Gln	Ile	Thr	Met	Glu	Lys																	
					185						190						195	200														
acg	ccc	agt	tac	ttc	gtc	acg	cgg	gag	gcc	ccc	gcg	cgc	atc	tgc	gcc	978																
Thr	Pro	Ser	Tyr	Phe	Val	Thr	Arg	Glu	Ala	Pro	Ala	Arg	Ile	Ser	Ala																	
					205						210						215															
atg	tcc	aag	gac	acc	aag	ctc	atc	gtg	gtg	gtg	cgg	gac	ccg	gtg	acc	1026																
Met	Ser	Lys	Asp	Thr	Lys	Leu	Ile	Val	Val	Val	Arg	Asp	Pro	Val	Thr																	
					220						225						230															

agg gcc atc tcg gac tac acg cag acg ctg tcc aag cgg ccc gac atc	1074
Arg Ala Ile Ser Asp Tyr Thr Gln Thr Leu Ser Lys Arg Pro Asp Ile	
235 240 245	
ccc acc ttc gag agc ttg acg ttc aaa aac agg aca gcg ggc ctc atc	1122
Pro Thr Phe Glu Ser Leu Thr Phe Lys Asn Arg Thr Ala Gly Leu Ile	
250 255 260	
gac acg tcg tgg agc gcc atc cag atc ggc atc tac gcc aag cac ctg	1170
Asp Thr Ser Trp Ser Ala Ile Gln Ile Gly Ile Tyr Ala Lys His Leu	
265 270 275 280	
gag cac tgg ctg cgc cac ttc ccc atc cgc cag atg ctc ttc gtg agc	1218
Glu His Trp Leu Arg His Phe Pro Ile Arg Gln Met Leu Phe Val Ser	
285 290 295	
ggc gag cgg ctc atc agc gac ccg gcc ggg gag ctg ggc cgc gtg caa	1266
Gly Glu Arg Leu Ile Ser Asp Pro Ala Gly Glu Leu Gly Arg Val Gln	
300 305 310	
gac ttc ctg ggc ctc aag agg atc atc acg gac aag cac ttc tac ttc	1314
Asp Phe Leu Gly Leu Lys Arg Ile Ile Thr Asp Lys His Phe Tyr Phe	
315 320 325	
aac aag acc aag ggc ttc ccc tgc ctg aag aag gcg gag ggc agc agc	1362
Asn Lys Thr Lys Gly Phe Pro Cys Leu Lys Lys Ala Glu Gly Ser Ser	
330 335 340	
cgg ccc cat tgc ctg ggc aag acc aag ggc agg acc cat cct gag atc	1410
Arg Pro His Cys Leu Gly Lys Thr Lys Gly Arg Thr His Pro Glu Ile	
345 350 355 360	
gac cgc gag gtg gtg cgc agg ctg cgc gag ttc tac cgg cct ttc aac	1458
Asp Arg Glu Val Val Arg Arg Leu Arg Glu Phe Tyr Arg Pro Phe Asn	
365 370 375	
ctc aag ttc tac cag atg acc ggg cac gac ttt ggc tgg gat	1500
Leu Lys Phe Tyr Gln Met Thr Gly His Asp Phe Gly Trp Asp	
380 385 390	
tgagcagacc cgggctatgt accttaccca cgtggcttat ctattgacag agattatatg	1560
tatgtaaaaat gtacagaaat ctattttata ataatttatt tttaattcat aagcaattaa	1620
ttcactaagc tgcctagcca cactcttttag agagtttagct tcataatctg ttaacattcc	1680
aaagtgttta actctagtat ttcgttttct tcttcacaat tgatgggtgct tctattttttt	1740
cttctcccct acctgttata tttaaaacaa agaaaagcac aacttgagat ttttgttggt	1800
acgggtattc agccttcagt caccgtctga gttctccagt tgctgcctcc ttgtcttgct	1860
ttgggtctcc cattccagct tcctgtctc ttctgcctg tgtacctcgt aggaacgctg	1920
agctgcctca acagggctgt attctgaagg gcaggcctca tgcagcagcc tccttgcaga	1980

tgtggtgtcc cgtccaatga tgtagcctga aagccacagc cctaggggttc tg

2032

<210> 10

<211> 390

<212> PRT

<213> Homo sapiens

<400> 10

Met Gly Gln Arg Leu Ser Gly Gly Arg Ser Cys Leu Asp Val Pro Gly  
1 5 10 15

Arg Leu Leu Pro Gln Pro Pro Pro Pro Pro Pro Pro Val Arg Arg Lys  
20 25 30

Leu Ala Leu Leu Phe Ala Met Leu Cys Val Trp Leu Tyr Met Phe Leu  
35 40 45

Tyr Ser Cys Ala Gly Ser Cys Ala Ala Ala Pro Gly Leu Leu Leu Leu  
50 55 60

Gly Ser Gly Ser Arg Ala Ala His Asp Pro Pro Ala Leu Ala Thr Ala  
65 70 75 80

Pro Asp Gly Thr Pro Pro Arg Leu Pro Phe Arg Ala Pro Pro Ala Thr  
85 90 95

Pro Leu Ala Ser Gly Lys Glu Met Ala Glu Gly Ala Ala Ser Pro Glu  
100 105 110

Glu Gln Ser Pro Glu Val Pro Asp Ser Pro Ser Pro Ile Ser Ser Phe  
115 120 125

Phe Ser Gly Ser Gly Ser Lys Gln Leu Pro Gln Ala Ile Ile Ile Gly  
130 135 140

Val Lys Lys Gly Gly Thr Arg Ala Leu Leu Glu Phe Leu Arg Val His  
145 150 155 160

Pro Asp Val Arg Ala Val Gly Ala Glu Pro His Phe Phe Asp Arg Ser  
165 170 175

Tyr Asp Lys Gly Leu Ala Trp Tyr Arg Asp Leu Met Pro Arg Thr Leu  
180 185 190

Asp Gly Gln Ile Thr Met Glu Lys Thr Pro Ser Tyr Phe Val Thr Arg  
195 200 205

Glu Ala Pro Ala Arg Ile Ser Ala Met Ser Lys Asp Thr Lys Leu Ile  
210 215 220

Val Val Val Arg Asp Pro Val Thr Arg Ala Ile Ser Asp Tyr Thr Gln  
225 230 235 240

Thr Leu Ser Lys Arg Pro Asp Ile Pro Thr Phe Glu Ser Leu Thr Phe  
245 250 255

004240" 2924360

Lys Asn Arg Thr Ala Gly Leu Ile Asp Thr Ser Trp Ser Ala Ile Gln  
 260 265 270

Ile Gly Ile Tyr Ala Lys His Leu Glu His Trp Leu Arg His Phe Pro  
 275 280 285

Ile Arg Gln Met Leu Phe Val Ser Gly Glu Arg Leu Ile Ser Asp Pro  
 290 295 300

Ala Gly Glu Leu Gly Arg Val Gln Asp Phe Leu Gly Leu Lys Arg Ile  
 305 310 315 320

Ile Thr Asp Lys His Phe Tyr Phe Asn Lys Thr Lys Gly Phe Pro Cys  
 325 330 335

Leu Lys Lys Ala Glu Gly Ser Ser Arg Pro His Cys Leu Gly Lys Thr  
 340 345 350

Lys Gly Arg Thr His Pro Glu Ile Asp Arg Glu Val Val Arg Arg Leu  
 355 360 365

Arg Glu Phe Tyr Arg Pro Phe Asn Leu Lys Phe Tyr Gln Met Thr Gly  
 370 375 380

His Asp Phe Gly Trp Asp  
 385 390

<210> 11  
 <211> 3658  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (847) .. (2214)

<220>  
 <223> Predicted human 3-OST-4 hnRNA

<400> 11  
 gaggatatcc cgggcgagag aaggagggt cggggatggg ctgagttgga gtcccagagg 60  
 aaaagcggaa gcgagagctt cgtcacccgc tgtcttccag ctcccgggtgc gcggcaccgg 120  
 aggcaggcgt tgggctttac ctctctaaaa gtactggggc aaaggaatgg agaacacggc 180  
 gtcccagact cccaagggag gggagtaaac gaggtggggg ggggaacacc ccaagtgcgt 240  
 gcgtgctggg gggctggggg gcacgatctc cgttctcccg ggtgccccag ccctagcgca 300  
 cgctccgct ccccgcccc ctctgcaggc gcgcgcgagg cgacaccccc ttccctcggc 360  
 ggcgcggggc gcgcgcccgg cccctctctc ctcccctcgg cgctctctct ctctcccggc 420  
 agaaagttag cagcggggaa ggaactctgg gctgcaacag cgcgcggcgg cggcggcaga 480



```

ggctgaagca gaagccgcgg cggagccggg gaagcggggg cgctgcagac ggagcaggtg 540
ccgccggcgg gtccgcgcgc cccctcggt ccccttgctt gaggctgagg ggggggcggt 600
ggtggggggg ccactcggac tcggcgggca gcgtggggcg gggggccatg cggccgggct 660
ccccctggc gcagcgggac agcggccagg gccgggggcg cagcggcgtc gcttcatgca 720
gccggggcgg ctgggcagcg gcggcggcgg cggcggcggc ggcggcgggg gcggcggctg 780
aaaccatgtc cgggcagcgc cgggggctgc cgccgcgcgc gccgccgcgc cgagccggga 840

gccgcg atg gcc cgg tgg ccc gca cct cct ccg cct ccg cct ccg cct 888
      Met Ala Arg Trp Pro Ala Pro Pro Pro Pro Pro Pro Pro Pro
        1             5             10

cca cct ctg gcc gcg ccg ccg ccg ccc ggc gcc tct gct aag ggg ccg 936
Pro Pro Leu Ala Ala Pro Pro Pro Pro Gly Ala Ser Ala Lys Gly Pro
  15             20             25             30

ccg gcg cgc aag ctg ctt ttt atg tgc acc ttg tcc ctg tct gtc acc 984
Pro Ala Arg Lys Leu Leu Phe Met Cys Thr Leu Ser Leu Ser Val Thr
        35             40             45

tac ctg tgc tac agc ctc ctg ggc ggc tgc ggc tcc ctg caa ttc cct 1032
Tyr Leu Cys Tyr Ser Leu Leu Gly Gly Ser Gly Ser Leu Gln Phe Pro
        50             55             60

ctg gcg ctg cag gag tgc ccg ggc gcc gcc gcc gag ccc ccg ccg agc 1080
Leu Ala Leu Gln Glu Ser Pro Gly Ala Ala Ala Glu Pro Pro Pro Ser
        65             70             75

ccg ccg cca ccc tct ctg ctg cct acc ccc gtg cgc ctc ggc gcc ccc 1128
Pro Pro Pro Pro Ser Leu Leu Pro Thr Pro Val Arg Leu Gly Ala Pro
        80             85             90

tcg cag ccg ccc gcg ccg ccg ccg ctg gac aac gcg agc cac ggg gag 1176
Ser Gln Pro Pro Ala Pro Pro Pro Leu Asp Asn Ala Ser His Gly Glu
        95             100             105             110

ccg ccc gag ccc cca gag cag cca gcc gcc ccc ggg acc gac ggc tgg 1224
Pro Pro Glu Pro Pro Glu Gln Pro Ala Ala Pro Gly Thr Asp Gly Trp
        115             120             125

ggg ctg ccg agc ggc ggc gga ggc gcc cgg gac gcc tgg ctc ccg acc 1272
Gly Leu Pro Ser Gly Gly Gly Gly Ala Arg Asp Ala Trp Leu Arg Thr
        130             135             140

ccg ctg gcc ccc agc gag atg atc acg gct cag agc gcg ctg ccg gag 1320
Pro Leu Ala Pro Ser Glu Met Ile Thr Ala Gln Ser Ala Leu Pro Glu
        145             150             155

agg gaa gcg cag gag tcc agc acc acc gac gag gat ctc gca ggc ccg 1368
Arg Glu Ala Gln Glu Ser Ser Thr Thr Asp Glu Asp Leu Ala Gly Arg
        160             165             170

aga gcg gcc aac ggg agc agc gag agg ggc ggc gcc gtc agc acc ccc 1416

```

Arg	Ala	Ala	Asn	Gly	Ser	Ser	Glu	Arg	Gly	Gly	Ala	Val	Ser	Thr	Pro	
175					180					185					190	
gac	tat	ggg	gag	aag	aag	ctg	cca	cag	gcg	ctc	atc	atc	ggg	gtc	aag	1464
Asp	Tyr	Gly	Glu	Lys	Lys	Leu	Pro	Gln	Ala	Leu	Ile	Ile	Gly	Val	Lys	
				195					200					205		
aaa	gga	ggg	acc	cgc	gcg	ctg	ctg	gag	gcg	atc	cgc	gtg	cac	ccg	gac	1512
Lys	Gly	Gly	Thr	Arg	Ala	Leu	Leu	Glu	Ala	Ile	Arg	Val	His	Pro	Asp	
			210					215					220			
gtg	cgg	gcg	gtg	ggc	gta	gag	ccg	cac	ttc	ttc	gac	agg	aac	tac	gaa	1560
Val	Arg	Ala	Val	Gly	Val	Glu	Pro	His	Phe	Phe	Asp	Arg	Asn	Tyr	Glu	
		225					230					235				
aag	ggg	ttg	gag	tgg	tac	aga	aat	gtg	atg	ccc	aag	act	ttg	gat	ggg	1608
Lys	Gly	Leu	Glu	Trp	Tyr	Arg	Asn	Val	Met	Pro	Lys	Thr	Leu	Asp	Gly	
	240					245					250					
caa	ata	acc	atg	gag	aag	act	cca	agt	tac	ttt	gtg	aca	aat	gag	gct	1656
Gln	Ile	Thr	Met	Glu	Lys	Thr	Pro	Ser	Tyr	Phe	Val	Thr	Asn	Glu	Ala	
255					260					265					270	
ccc	aag	cgc	att	cac	tcc	atg	gcc	aag	gac	atc	aaa	ctg	att	gtg	gtg	1704
Pro	Lys	Arg	Ile	His	Ser	Met	Ala	Lys	Asp	Ile	Lys	Leu	Ile	Val	Val	
				275					280					285		
gtg	aga	aac	ccc	gtg	acc	agg	gcc	atc	tct	gac	tac	acg	cag	aca	ctg	1752
Val	Arg	Asn	Pro	Val	Thr	Arg	Ala	Ile	Ser	Asp	Tyr	Thr	Gln	Thr	Leu	
			290					295					300			
tca	aag	aaa	ccc	gag	atc	ccc	acc	ttt	gag	gtg	ctg	gcc	ttc	aaa	aac	1800
Ser	Lys	Lys	Pro	Glu	Ile	Pro	Thr	Phe	Glu	Val	Leu	Ala	Phe	Lys	Asn	
		305					310					315				
cgg	acc	ctc	ggg	ctg	atc	gat	gct	tcc	tgg	agt	gcc	att	cga	ata	ggg	1848
Arg	Thr	Leu	Gly	Leu	Ile	Asp	Ala	Ser	Trp	Ser	Ala	Ile	Arg	Ile	Gly	
	320					325					330					
atc	tat	gcg	ctg	cat	ctg	gaa	aac	tgg	ctc	cag	tat	ttc	ccc	ctc	tcc	1896
Ile	Tyr	Ala	Leu	His	Leu	Glu	Asn	Trp	Leu	Gln	Tyr	Phe	Pro	Leu	Ser	
335					340					345					350	
cag	atc	ctc	ttt	gtc	agt	ggc	gag	cga	ctc	att	gtg	gac	ccc	gcc	ggg	1944
Gln	Ile	Leu	Phe	Val	Ser	Gly	Glu	Arg	Leu	Ile	Val	Asp	Pro	Ala	Gly	
				355					360					365		
gaa	atg	gcc	aaa	gta	cag	gat	ttt	cta	ggc	ctc	aaa	cgt	gtt	gtg	act	1992
Glu	Met	Ala	Lys	Val	Gln	Asp	Phe	Leu	Gly	Leu	Lys	Arg	Val	Val	Thr	
			370					375					380			
aag	aag	cat	ttc	tat	ttc	aac	aaa	acc	aag	ggg	ttc	cct	tgc	cta	aag	2040
Lys	Lys	His	Phe	Tyr	Phe	Asn	Lys	Thr	Lys	Gly	Phe	Pro	Cys	Leu	Lys	
		385					390					395				
aag	cca	gaa	gac	agc	agt	gcc	ccg	agg	tgc	tta	ggc	aag	agc	aaa	ggc	2088
Lys	Pro	Glu	Asp	Ser	Ser	Ala	Pro	Arg	Cys	Leu	Gly	Lys	Ser	Lys	Gly	

400

405

410

cgg act cat cct cgc att gac cca gat gtc atc cac aga ctg agg aaa 2136  
Arg Thr His Pro Arg Ile Asp Pro Asp Val Ile His Arg Leu Arg Lys  
415 420 425 430

ttc tac aaa ccc ttc aac ttg atg ttt tac caa atg act ggt caa gat 2184  
Phe Tyr Lys Pro Phe Asn Leu Met Phe Tyr Gln Met Thr Gly Gln Asp  
435 440 445

ttt cag tgg gaa cag gaa gag ggt gat aaa tgaggctaga gaggcagagg 2234  
Phe Gln Trp Glu Gln Glu Glu Gly Asp Lys  
450 455

aaggctagtc aataagctaa ggaggctcct tgcctgagtc cttgaatacc ccagcttctg 2294

cagcttcact tgctggagtg ccaagtagat ctccctcctcc ttcattgcagc caggattgcc 2354

tccagtgcctg ttagcttagg caaacagggtg gatcccatgg catcccatg gaggaaccag 2414

gcccattctgg gcagcagcat ctggttgacc agatggccac cagaaccac tggtcattct 2474

tatcttctgc tagttaatat agcctgaaga cagaggataa atagttgtca atgtcagaga 2534

cagtgcctatt aatgtatatg tgagcgacaa aaaagggtctg ctttatagggt gttctcactc 2594

tagcttgggg agcccagggt tctagccctg tatctgtcat gggcacctgc tgtctaaacc 2654

tctgcttggg cttctcccca gaatgcactt tgtggctgag tgctccagga ctccctaggga 2714

gcaagctcct ccctctaagg tgtttctagt cttctcttta aagggtctcat cccacaaccc 2774

ctgacttcct ccctcccccac atcatgaagg cagaggcatg cacattcctc actgaaaaag 2834

aaaacacaca cccaccacaca cacacacaca cagaagaaaa tgaaagctga cacacctcga 2894

agccttcttt ccaagagccc tctaaatggg gttgggtctc actcttcatg agtatcctgg 2954

gttgtgcaga agcttagcat atgcccttgt gttcggatca ggcccacagg gctgctcaaa 3014

gagtagagta attgtaaccg aggtcagagc tctgggggttg gcagagatga gtggccatat 3074

ctgggggttaa aagaagaaat cctgtcctct tgggtgggagg ttaccttacc tgaagacat 3134

ctctcccaag cactgtagtt ctgagcatgt ttttgggggtg gactctgtcc cctagggtcc 3194

ctagaagggtc aaagaccaga gagttgacaa gtctgttatt aggaataatc cttagccatg 3254

taatggagaa aggagcagtc agcattcttc caatttgccc caccaccacc tcctcggggt 3314

tcattttctc tatttagaga tggcagagag tgaggtagtg gcgagaaagc tgactccatt 3374

catcagatcc agtttatgag ggttgggggt gagcaagggtc tgtctgcaga aacccccatc 3434

aagagctgct gaatgaagtg tcccttccca tcagtttgat tcaattaaaa tgcattcattt 3494

gacataaagc acttgttcac agatctccaa aaccagggaat tgttctagta aaactggaaa 3554

tttgtatgag tgggggggagt taaatctgtt cagctgttat taaactgtca tttctcccg 3614

taaatgaaaa ccgtgttgtt ataaagctta atgcaacctg atta 3658

<210> 12

<211> 456

<212> PRT

<213> Homo sapiens

<400> 12

Met Ala Arg Trp Pro Ala Pro Pro Pro Pro Pro Pro Pro Pro Pro Pro  
1 5 10 15

Leu Ala Ala Pro Pro Pro Pro Gly Ala Ser Ala Lys Gly Pro Pro Ala  
20 25 30

Arg Lys Leu Leu Phe Met Cys Thr Leu Ser Leu Ser Val Thr Tyr Leu  
35 40 45

Cys Tyr Ser Leu Leu Gly Gly Ser Gly Ser Leu Gln Phe Pro Leu Ala  
50 55 60

Leu Gln Glu Ser Pro Gly Ala Ala Ala Glu Pro Pro Pro Ser Pro Pro  
65 70 75 80

Pro Pro Ser Leu Leu Pro Thr Pro Val Arg Leu Gly Ala Pro Ser Gln  
85 90 95

Pro Pro Ala Pro Pro Pro Leu Asp Asn Ala Ser His Gly Glu Pro Pro  
100 105 110

Glu Pro Pro Glu Gln Pro Ala Ala Pro Gly Thr Asp Gly Trp Gly Leu  
115 120 125

Pro Ser Gly Gly Gly Gly Ala Arg Asp Ala Trp Leu Arg Thr Pro Leu  
130 135 140

Ala Pro Ser Glu Met Ile Thr Ala Gln Ser Ala Leu Pro Glu Arg Glu  
145 150 155 160

Ala Gln Glu Ser Ser Thr Thr Asp Glu Asp Leu Ala Gly Arg Arg Ala  
165 170 175

Ala Asn Gly Ser Ser Glu Arg Gly Gly Ala Val Ser Thr Pro Asp Tyr  
180 185 190

Gly Glu Lys Lys Leu Pro Gln Ala Leu Ile Ile Gly Val Lys Lys Gly  
195 200 205

Gly Thr Arg Ala Leu Leu Glu Ala Ile Arg Val His Pro Asp Val Arg  
210 215 220

Ala Val Gly Val Glu Pro His Phe Phe Asp Arg Asn Tyr Glu Lys Gly  
225 230 235 240

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Leu Glu Trp Tyr Arg Asn Val Met Pro Lys Thr Leu Asp Gly Gln Ile  
 245 250 255  
 Thr Met Glu Lys Thr Pro Ser Tyr Phe Val Thr Asn Glu Ala Pro Lys  
 260 265 270  
 Arg Ile His Ser Met Ala Lys Asp Ile Lys Leu Ile Val Val Val Arg  
 275 280 285  
 Asn Pro Val Thr Arg Ala Ile Ser Asp Tyr Thr Gln Thr Leu Ser Lys  
 290 295 300  
 Lys Pro Glu Ile Pro Thr Phe Glu Val Leu Ala Phe Lys Asn Arg Thr  
 305 310 315 320  
 Leu Gly Leu Ile Asp Ala Ser Trp Ser Ala Ile Arg Ile Gly Ile Tyr  
 325 330 335  
 Ala Leu His Leu Glu Asn Trp Leu Gln Tyr Phe Pro Leu Ser Gln Ile  
 340 345 350  
 Leu Phe Val Ser Gly Glu Arg Leu Ile Val Asp Pro Ala Gly Glu Met  
 355 360 365  
 Ala Lys Val Gln Asp Phe Leu Gly Leu Lys Arg Val Val Thr Lys Lys  
 370 375 380  
 His Phe Tyr Phe Asn Lys Thr Lys Gly Phe Pro Cys Leu Lys Lys Pro  
 385 390 395 400  
 Glu Asp Ser Ser Ala Pro Arg Cys Leu Gly Lys Ser Lys Gly Arg Thr  
 405 410 415  
 His Pro Arg Ile Asp Pro Asp Val Ile His Arg Leu Arg Lys Phe Tyr  
 420 425 430  
 Lys Pro Phe Asn Leu Met Phe Tyr Gln Met Thr Gly Gln Asp Phe Gln  
 435 440 445  
 Trp Glu Gln Glu Glu Gly Asp Lys  
 450 455

<210> 13

<211> 284

<212> PRT

<213> Homo sapiens

<220>

<223> human NST-1 (aa 599 to 882)

<400> 13

Lys Thr Cys Asp Arg Phe Pro Lys Leu Leu Ile Ile Gly Pro Gln Lys  
 1 5 10 15

Thr Gly Thr Thr Ala Leu Tyr Leu Phe Leu Gly Met His Pro Asp Leu  
 20 25 30

Ser Ser Asn Tyr Pro Ser Ser Glu Thr Phe Glu Glu Ile Gln Phe Phe  
35 40 45

Asn Gly His Asn Tyr His Lys Gly Ile Asp Trp Tyr Met Glu Phe Phe  
50 55 60

Pro Ile Pro Ser Asn Thr Thr Ser Asp Phe Tyr Phe Glu Lys Ser Ala  
65 70 75 80

Asn Tyr Phe Asp Ser Glu Val Ala Pro Arg Arg Ala Ala Ala Leu Leu  
85 90 95

Pro Lys Ala Lys Val Leu Thr Ile Leu Ile Asn Pro Ala Asp Arg Ala  
100 105 110

Tyr Ser Trp Tyr Gln His Gln Arg Ala His Asp Asp Pro Val Ala Leu  
115 120 125

Lys Tyr Thr Phe His Glu Val Ile Thr Ala Gly Ser Asp Ala Ser Ser  
130 135 140

Lys Leu Arg Ala Leu Gln Asn Arg Cys Leu Val Pro Gly Trp Tyr Ala  
145 150 155 160

Thr His Ile Glu Arg Trp Leu Ser Ala Tyr His Ala Asn Gln Ile Leu  
165 170 175

Val Leu Asp Gly Lys Leu Leu Arg Thr Glu Pro Ala Lys Val Met Asp  
180 185 190

Met Val Gln Lys Phe Leu Gly Val Thr Asn Thr Ile Asp Tyr His Lys  
195 200 205

Thr Leu Ala Phe Asp Pro Lys Lys Gly Phe Trp Cys Gln Leu Leu Glu  
210 215 220

Gly Gly Lys Thr Lys Cys Leu Gly Lys Ser Lys Gly Arg Lys Tyr Pro  
225 230 235 240

Glu Met Asp Leu Asp Ser Arg Ala Phe Leu Lys Asp Tyr Tyr Arg Asp  
245 250 255

His Asn Ile Glu Leu Ser Lys Leu Leu Tyr Lys Met Gly Gln Thr Leu  
260 265 270

Pro Thr Trp Leu Arg Glu Asp Leu Gln Asn Thr Arg  
275 280

<210> 14

<211> 286

<212> PRT

<213> Homo sapiens

<220>

<223> human NST-2 (aa 598 to 883)

<400> 14

Lys	Thr	Cys	Asp	Arg	Leu	Pro	Lys	Phe	Leu	Ile	Val	Gly	Pro	Gln	Lys	1	5	10	15
Thr	Gly	Thr	Thr	Ala	Ile	His	Phe	Phe	Leu	Ser	Leu	His	Pro	Ala	Val	20	25	30	
Thr	Ser	Ser	Phe	Pro	Ser	Pro	Ser	Thr	Phe	Glu	Glu	Ile	Gln	Phe	Phe	35	40	45	
Asn	Ser	Pro	Asn	Tyr	His	Lys	Gly	Ile	Asp	Trp	Tyr	Met	Asp	Phe	Phe	50	55	60	
Pro	Val	Pro	Ser	Asn	Ala	Ser	Thr	Asp	Phe	Leu	Phe	Glu	Lys	Ser	Ala	65	70	75	80
Thr	Tyr	Phe	Asp	Ser	Glu	Val	Val	Pro	Arg	Arg	Gly	Ala	Ala	Leu	Leu	85	90	95	
Pro	Arg	Ala	Lys	Ile	Ile	Thr	Val	Leu	Thr	Asn	Pro	Ala	Asp	Arg	Ala	100	105	110	
Tyr	Ser	Trp	Tyr	Gln	His	Gln	Arg	Ala	His	Gly	Asp	Pro	Val	Ala	Leu	115	120	125	
Asn	Tyr	Thr	Phe	Tyr	Gln	Val	Ile	Ser	Ala	Ser	Ser	Gln	Thr	Pro	Leu	130	135	140	
Ala	Leu	Arg	Ser	Leu	Gln	Asn	Arg	Cys	Leu	Val	Pro	Gly	Tyr	Tyr	Ser	145	150	155	160
Thr	His	Leu	Gln	Arg	Trp	Leu	Thr	Tyr	Tyr	Pro	Ser	Gly	Gln	Leu	Leu	165	170	175	
Ile	Val	Asp	Gly	Gln	Glu	Leu	Arg	Thr	Asn	Pro	Ala	Ala	Ser	Met	Glu	180	185	190	
Ser	Ile	Gln	Lys	Phe	Leu	Gly	Ile	Thr	Pro	Phe	Leu	Asn	Tyr	Thr	Arg	195	200	205	
Thr	Leu	Arg	Phe	Asp	Asp	Asp	Lys	Gly	Phe	Trp	Cys	Gln	Gly	Leu	Glu	210	215	220	
Gly	Gly	Lys	Thr	Arg	Cys	Leu	Gly	Arg	Ser	Lys	Gly	Arg	Arg	Tyr	Pro	225	230	235	240
Asp	Met	Asp	Thr	Glu	Ser	Arg	Leu	Phe	Leu	Thr	Asp	Phe	Phe	Arg	Asn	245	250	255	
His	Asn	Leu	Glu	Leu	Ser	Lys	Leu	Leu	Ser	Arg	Leu	Gly	Gln	Pro	Val	260	265	270	
Pro	Ser	Trp	Leu	Arg	Glu	Glu	Leu	Gln	His	Ser	Ser	Leu	Gly	275	280	285			

<210> 15  
 <211> 291  
 <212> PRT  
 <213> Caenorhabditis elegans

<220>  
 <223> putative C. elegans 3-OST

<400> 15

Met Lys Tyr Arg Leu Leu Leu Ile Leu His Leu Ile Asp Leu Ile Ser  
 1 5 10 15

Cys Gly Val Ile Pro Asn Thr Ser Lys Lys Arg Phe Pro Asp Ala Ile  
 20 25 30

Ile Val Gly Val Lys Lys Ser Gly Thr Arg Ala Leu Leu Glu Phe Leu  
 35 40 45

Arg Val Asn Pro Leu Ile Lys Ala Pro Gly Pro Glu Val His Phe Phe  
 50 55 60

Asp Lys Asn Phe Asn Lys Gly Leu Glu Trp Tyr Arg Glu Gln Met Pro  
 65 70 75 80

Glu Thr Lys Phe Gly Glu Val Thr Ile Glu Lys Ser Pro Ala Tyr Phe  
 85 90 95

His Ser Lys Met Ala Pro Glu Arg Ile Lys Ser Leu Asn Pro Asn Thr  
 100 105 110

Lys Ile Ile Ile Val Val Arg Asp Pro Val Thr Arg Ala Ile Ser Asp  
 115 120 125

Tyr Thr Gln Ser Ser Ser Lys Arg Lys Arg Val Gly Leu Met Pro Ser  
 130 135 140

Phe Glu Thr Met Ala Val Gly Asn Cys Ala Asn Trp Leu Arg Thr Asn  
 145 150 155 160

Cys Thr Thr Lys Thr Arg Gly Val Asn Ala Gly Trp Gly Ala Ile Arg  
 165 170 175

Ile Gly Val Tyr His Lys His Met Lys Arg Trp Leu Asp His Phe Pro  
 180 185 190

Ile Glu Asn Ile His Ile Val Asp Gly Glu Lys Leu Ile Ser Asn Pro  
 195 200 205

Ala Asp Glu Ile Ser Ala Thr Glu Lys Phe Leu Gly Leu Lys Pro Val  
 210 215 220

Ala Lys Pro Glu Lys Phe Gly Val Asp Pro Ile Lys Lys Phe Pro Cys  
 225 230 235 240

Ile Lys Asn Glu Asp Gly Lys Leu His Cys Leu Gly Lys Thr Lys Gly  
 245 250 255



Arg His His Pro Asp Val Glu Pro Ser Val Leu Lys Thr Leu Arg Glu  
260 265 270

Phe Tyr Gly Pro Glu Asn Lys Lys Phe Tyr Gln Met Ile Asn His Trp  
275 280 285

Phe Asp Trp  
290

<210> 16

<211> 4045

<212> DNA

<213> Homo sapiens

<220>

<223> 3-OST-4 5' promoter/exon

<400> 16

gaattctgtg ggtgttgga ggggagacag aaaactatct tccatcgagt cttcggatcc 60  
attgggaatg cctggatgac gtcagagttc gccctgtgta ggtagctccc acttttcatt 120  
gtaggtttct caaggacttg ctctagaaa aagcgtggct caaaagtaga taaaaaatag 180  
gcaactgcct aagtgtgaaa ttacaaaagt tcctctccaa aaaagcccgc ctctcccta 240  
tcacttgtag gcctgacatt ttaccaaagg ggctctatct tttcaagagt ttgttattaa 300  
agcgtgacta tttgaggatt ggaggcaaaa gggatactga gaaatgtcct tactagcagt 360  
gtcaaggcaa gtgacataaa tgtgtggggg ggcaacttgt atgagcactg tgaaaacggc 420  
agcatgttca ctctacttct cagctctgac tgaggggctc aaagttcagg atctgctgat 480  
ttttcaacag taacgtctc tccaagggtgt tttttttttt tccttttttg ggaaagcccc 540  
cagtttaaac tattgcagcc agtttacatt tcttaatgtc actgtgctgg ccacattcag 600  
agctccattt gccaccatcg gttttgatac ctttttacca aaacctttcg aaatttgaga 660  
gcccatcttt agtaaaactg ggcattggagc agattcgttt ggattgctga gaggggagat 720  
agaaaagttt ggggtgctagg caggaactgc aaggaggacc tgggcatat gccagacatc 780  
tagtgccctg gccttgaaag ggagactggt cgtgacaag gcaatatctg ttgcaacca 840  
ggcttcttag atgaccacct tggatcatgg ctcgagcac agggagggct gggcagtgtc 900  
tgtgtttctc tccgttccag ttggcccctt cccattgaca ttacagtaat gcagttgtgt 960  
gctgtttgaa aaagcatccc tagttacaca gaatgattta caggacacca gactctgcat 1020  
ttcagaggtc tccagtgtac cataaaaaat atattataaa agaataatct ttatctgaac 1080  
taaagctgca gtgaaggaaa ctctgtgtcca gctgagagca gcagtgtgct tttgttcaact 1140

cagggaaaaag	tccgtgttct	ttatcttatt	tgattaacat	tttttttctc	tttggcacta	1200
ggatatctcgt	taatatTTtag	acattatata	catttttcttt	caactggttt	tcctattacg	1260
tgatcaaaca	aaaccagaga	tgccagccac	agcagccaac	aagggaaaag	cagccccgtc	1320
aggcaccacag	ctgtggctgg	ggggcaaatg	gtactcacta	gagccacccc	caggaggcac	1380
ctggcagagc	tctgtgcaga	gccagccccg	gttgcagaaa	gctgagtttg	ttggagtgcc	1440
tcagttgatc	actctgtctc	tttctcccat	ttccctcact	tccctgagca	aaatgcaaca	1500
ggaagcaaag	tctagttgtg	aatcttccaa	agccttctga	tgtttaccat	gttcccccag	1560
gagaggggagg	tgaggggtgg	agatctctct	gcaaagaaaa	tacacttaaa	aaatttcagc	1620
gagccgatgc	acagacaccc	agcaaccacg	cttgtctccg	cttattaggt	gttcagagcg	1680
acagtgggtcc	cacactatTT	cagtccagga	aaccatgaac	tccgttagtg	gcaatgcccc	1740
cgaagaggcg	caggtgtgtg	cacctgtgat	taagggtgtc	gaggaggggc	agcctcatct	1800
cttgaagcag	aaagtgttgt	cacctgggtga	tgggacagag	ggaaaagctc	tggggctggg	1860
aaacctgggg	gcttgtgtca	aagctccacc	catcaggagc	ttcaagagaa	gatggggggc	1920
ggggggcggt	ggctggaaaag	atggaagttg	ggatgggaaa	gcggttgtag	aaaaggattc	1980
actcctggac	cgaaggcagg	aggatatccc	gggcgagaga	agggagggtc	ggggatgggc	2040
tgagttggag	tcccagagga	aaagcggaag	cgagagcttc	gtcacccgct	gtcttccagc	2100
tcccggtgcg	cggcaccgga	ggcaggcggt	gggcctttacc	tctctaaaag	tactggggca	2160
aaggaatgga	gaacacggcg	tcccgagctc	ccaagggagg	ggagtaaacg	aggtggggtg	2220
gggaacaccc	caagtgcgtg	cgtgctgggg	ggctgggggg	cacgatctcc	gttctcccg	2280
gtgccccagc	cctagcgcac	gcctccgctc	ccccgcccc	ttcgcaggcg	cgcgcgaggc	2340
gcacccccct	tccttcggcg	gcgcggggcg	cgcgcccggc	cccctcctcc	tcccctccgc	2400
gcctctcctc	tctcccgga	gaaagttagc	agcggggaaag	gaactctggg	ctgcaacagc	2460
gcgcggcggc	ggcggcagag	gctgaagcag	aagccgcggc	ggagccgggg	aagcgggggc	2520
gctgcagacg	gagcaggtgc	cgccggcggg	tccgcgcgcc	cccctcggtc	cccttgccctg	2580
aggctgaggg	gggggcggtg	gtgggggggc	cactcggact	cggcgggcag	cgtggggcgg	2640
ggggccatgc	ggccgggctc	ccccctggcg	cagcgggaca	gcggccaggg	ccgggggctc	2700
agcggcgctcg	cttcatgcag	ccggggcggc	tgggcagcgg	cggcggcggc	ggcggcgggc	2760
gcggcggggg	cggcggtga	aaccatgtcc	gggcagcgcc	gggggctgcc	gccgccggcg	2820
ccgccgccgc	gagccgggag	ccgcgatggc	ccggtggccc	gcacctctc	cgcctccgcc	2880

tccgcctcca cctctggcgc cgcgcgcgc gcccggcgc tctgctaagg ggccgcgcgc 2940  
 gcgcaagctg ctttttatgt gcaccttgtc cctgtctgtc acctacctgt gctacagcct 3000  
 cctgggcgcgc tcgggctccc tgcaattccc tctggcgctg caggagtgcg cgggcgcgcgc 3060  
 cgccgagccc ccgccgagcc cgccgccacc ctctctgctg cctacccccg tgcgcctcgc 3120  
 cgccccctcg cagccgcccgc cgccgcgcgc gctggacaac gcgagccacg gggagccgcgc 3180  
 cgagccccca gagcagccag ccgccccgcg gaccgacggc tgggggctgc cgagcggcgcg 3240  
 cggagggcgc cgggacgcct ggctccggac ccgctgggc cccagcgaga tgatcacggc 3300  
 tcagagcgcg ctgccggaga gggaagcgca ggagtccagc accaccgacg aggatctcgc 3360  
 aggccggaga gcggccaacg ggagcagcga gaggggcgcg gccgtcagca cccccgacta 3420  
 tggggagaag aagctgccac aggcgctcat catcggggtc aagaaaggag ggaccgcgcgc 3480  
 gctgctggag gcgatccgcg tgcaccgcga cgtgcgggcg gtgggcgtag agccgcactt 3540  
 cttcgacagg aactacgaaa aggggttgga gtggtacagg taggacctg ggctccgcgcg 3600  
 gctggtggag acgctggtgg gagacgcgga ggggaagccg cgctttcca cgcccttcga 3660  
 gcatccaggc accgtcccga gaggcccaag ccccgcgag ggctctgcaa accctggcgcg 3720  
 cgttgctcag ggggatcggc tgagagggct ggactccagc gaaaggtcac tttatttcag 3780  
 ggcgagggga ggaggtgtca ccctgccctg cctccgcgc tcctcatcca aggaggtgct 3840  
 gtctgaatct gccagctcc aagcctggga accccagcc ctctgctg ctgggtgttt 3900  
 ccgaaaccag gctcttgccg ggttctggga ttctgggcag aggactttga ggagtgcgac 3960  
 aggatggcta aattgactaa ggggatttga ggtcccctgg aatctcttaa aatcacccctc 4020  
 aaacgcattt gcgtggctgg aattc 4045